

ENGINEERING ENZYMES THROUGH GENETIC SELECTION**CROSS-REFERENCED TO RELATED APPLICATIONS**

This application claims benefit of and priority to US Provisional Patent Application No. 60/520,754 filed on November 17, 2003, US Provisional Patent Application No. 60/520,813, also filed on November 17, 2003, and US Provisional Patent Application No. 60/_____ filed on October 18, 2004, and where permissible, each of which is incorporated by reference in their entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

Aspects of the work described herein were supported in part by Grant No. DBI-0320786 award by the National Science Foundation. The US government may have certain rights in the disclosed subject matter.

1. Technical Field

Aspects of the present disclosure are generally directed to systems and methods for generating ligand-receptor pairs for transcriptional control by small molecules.

2. Related Art

Directed molecular evolution of enzymes is a developing field in the biotechnology industry and occurs through the single or repeated application of two steps: diversity/library generation followed by screening or selecting for function. The last several years have produced much progress in each of these areas. Techniques of diversity generation in the creation of libraries range from methods with no structure/function prejudice (error-prone PCR; mutator strains) to highly focused randomization based on structural information (site-directed mutagenesis; cassette mutagenesis). DNA recombination (DNA-shuffling, StEP, SCRATCHY, RACHITT, RDA-PCR) requires no structural information but works on the premise that Nature has already solved the problem of creating functional proteins from amino acids. By randomly recombining the genes for related proteins, new combinations of the different solutions are created which may be better than any of the original individual proteins. Structure-based approaches can be combined with other methods to generate greater diversity.

Advances have also been made in screening the generated libraries for proteins with desired properties. In a screen each protein in the library is

analyzed for function, which limits library size. In contrast, genetic selection evaluates entire libraries at once, in a highly parallel fashion, because only functional members of the library survive the selective pressure. In selection, nonfunctional members of the library are not individually evaluated. For screens, each variant must be individually assayed and the data evaluated, requiring more time and materials. In vivo genetic selection strategies enable the exhaustive analysis of protein libraries with up to about 10^{10} different members. The quoted throughputs are maximal values for industrial, robot driven laboratories. Realistically, experience indicates that an academic, individual investigator laboratory can achieve up to 10^4 samples/day for screening in yeast and 10^7 samples/day for genetic selection in yeast. In summary, genetic selection is generally preferable to screening not only because it is higher throughput, but also because it requires less time and materials.

With regard to selection, there are several common conventional selection strategies, such as i) antibiotic resistance, ii) substrate selected growth, where degradation of substrates provides elements essential for growth (such as C, N, P, and S), iii) auxotrophic complementation to restore metabolic function, and iv) phage display, which displays peptides or proteins on a virus surface and segregates them on the basis of binding affinity. Although powerful, these selection strategies are not general enough to apply to engineering enzymes for many interesting reactions. Conventional systems rely on screening techniques rather than selection techniques because selections are more difficult.

The generation of libraries has spawned many companies, in fact, spawned an industry. What has so far failed to be addressed is a general method of evaluating libraries (no matter how they are generated) through genetic selection. Accordingly there is a need for new compositions and methods for engineering polypeptides and rapidly identifying engineered polypeptides having desirable characteristics.

SUMMARY

Methods and compositions for selecting or screening transformed cells are provided. An exemplary method includes selecting transformed cells by introducing a first polynucleotide into a transformed cell unable to survive on selective media in the absence of a selection agent, wherein the transformed cell expresses a recombinant receptor polypeptide that activates transcription of a

second polynucleotide in response to interaction of the recombinant receptor polypeptide with a target substance, culturing the transformed cell on the selective media in the absence of the selection agent; and selecting the transformed cell that survives on the selective media in the absence of the selection agent.

Another aspect provides a method for selecting transformed cells by introducing a first polynucleotide into a transformed cell, wherein the transformed cell expresses a recombinant receptor polypeptide that activates transcription of a second polynucleotide in response to interaction of the recombinant receptor polypeptide with a target substance, culturing the transformed cell on the selective media in the presence of a first selection agent, and selecting the transformed cell that survives on the selective media in the absence of the selection agent, wherein the second polynucleotide encodes an enzyme that converts the first selective agent into a product toxic to the transformed cell.

Still another embodiment provides a cell including a recombinant nuclear receptor that induces transcription of a first polynucleotide in response to interaction with a target substance, and an adapter fusion protein comprising a human coactivator domain operably linked to an activation domain, wherein the adapter fusion protein enhances transcription of the first polynucleotide induced by the recombinant nuclear receptor.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows a schematic depicting an exemplary chemical complementation scheme. For selection, yeast strain PJ69-4A has the *ADE2* gene under the control of a Gal4 response element (Gal4RE). This strain is transformed with a plasmid expressing ACTR:GAD (manuscript submitted). Plasmids created through homologous recombination in PJ69-4A express a variant GBD:RXR. In media lacking adenine, yeast will grow only in the presence of a ligand that causes the RXR LBD to associate with ACTR and activate transcription of *ADE2*. For clarity, only one ACTR:GAD is depicted.

Figs. 2a-o are line graphs showing selection assay (SC -Ade -Trp -Leu + ligand) data for yeast growth in the presence of 9cRA (closed circles) and LG335 (open circles) for 43 hours.

Figs. 3a-o are line graphs showing screen assay (SC -Trp -Leu + ligand) data for β -galactosidase activity with *o*-Nitrophenyl β -D-galactopyranoside (ONPG) substrate in the presence of 9cRA (closed circles) and LG335 (open circles). Miller units normalize the change in absorbance at 405 nm for the change optical density at 630 nm, which reflects the number of cells per well.

Figs 4a and b are line graphs showing data from mammalian cell culture using a luciferase reporter with wtRXR (solid circle), I268A;I310S;F313A;L436F (solid dot), I268V;A272V;I310M;F313S;L436M (inverted triangle), I268A;I310M;F313A;L436T (gray square), I268V;A272V;I310L;F313M (upright triangle), or I268A;I310A;F313A;L436F (grey circle) in response to (a) 9cRA and LG335 (b). RLU = relative light units.

Figs. 5a-g are photographs of culture plates showing yeast transformed with both ACTR:GAD and GBD:RXR grow in the presence of various concentrations of 9cRA.

Figs. 6a-g are photographs of culture plates showing yeast transformed with both SRC-1:GAD and GBD:RXR grow in the presence of various concentrations of 9cRA.

Figs. 7a-f are photographs of culture plates showing negative selection of yeast transformed with both ACTR:GAD and GBD:RXR in the presence of various concentrations of 9cRA.

Figs. 8a-t are photographs of culture plates showing growth due to the indicated transformants of variant GBD:RXRs due to various concentrations of 9cRA.

Figs. 9a -e are schematics of exemplary embodiments for the selection of desired transformants.

Fig. 10 is a schematic of an exemplary embodiment for the selection of selective receptor modulators in transformants incorporating a human nuclear receptor coactivator fused to a repression domain.

Fig. 11 is a schematic of an exemplary embodiment for the selection of receptor antagonists.

Fig. 12 is a schematic of an exemplary embodiment for chemical complementation selection of transformants to obtain isotype or isoform selective receptor agonists.

Fig. 13 is a schematic of an exemplary embodiment for chemical complementation selection of transformants incorporating a nuclear receptor coactivator fused to an activation domain for the selection of receptor agonists.

Fig. 14 is a Ligplot depiction of hydrophobic interactions between the RXR LBD and 9cRA.

Figs. 15a-b show the structure of exemplary ligands used in chemical complementation of one embodiment.

Figs. 16a-b show schematics of exemplary methods for the construction of pGBDRXR:3stop (a) or an insert cassette library (b).

Figs. 17a-b are diagrams of exemplary constructs according to one embodiment of the present disclosure.

DETAILED DESCRIPTION

Methods and compositions for engineering proteins are provided, in particular, methods for engineering proteins that interact with a target compound. Embodiments of the disclosure combine chemical complementation with genetic selection to engineer proteins, polypeptides, enzymes, antibodies, adhesins, integrins, and the like. Typically, any protein or polypeptide that interacts with a small molecule can be engineered or modified using the disclosed methods and systems. Exemplary proteins include, but are not limited to enzymes, antibodies, cell surface receptors, polypeptides involved in signal transduction pathways, intracellular polypeptides, secreted polypeptides, and transmembrane polypeptides. In some embodiments, the polypeptides interact with a small molecule that is produced naturally. Representative naturally produced small molecules include but are not limited to, neurotransmitters, cAMP, cGMP, steroids, purines, pyrimidines, heterocyclic compounds, ATP, DAG, IP3, inositol, calcium ions, magnesium ions, vitamins, minerals, and combinations thereof. Some embodiments provide methods and systems for engineering proteins that distinguish between optical isomers of a target compound.

Other embodiments provide a more efficient mammalian model system in yeast for evaluating protein/ligand interactions, and can be utilized in an array of applications including but not limited to drug discovery. Nuclear receptors are implicated in diseases such as diabetes and various cancers. Agonists and antagonists for these nuclear receptors serve as drugs. With chemical

complementation, libraries of compounds can be screened as potential agonists, as described herein. In some embodiments, antagonists can be identified with negative chemical complementation. Chemical complementation can also be extended to identify isotype-selective agonists and antagonists and used for the discovery of selective receptor modulators (e.g., SERMs).

In addition to drug discovery, the increase in sensitivity of disclosed systems and methods also provides a method for engineering receptors to recognize small molecules. For example, libraries of engineered receptors can be transformed into yeast and plated onto media containing the target ligand.

These engineered receptors can be used for controlling transcription in mammalian cells, and potentially applied towards gene therapy. Furthermore, some embodiments of the disclosed system can give insight into the general mechanism for understanding the fundamentals of protein structure and function.

In summary, we have demonstrated that the addition of an adapter protein consisting of a human coactivator fused to a yeast transcriptional activator increases the sensitivity of chemical complementation with RXR 1000-fold, enhancing the system so that it is indistinguishable from activation by Gal4. Negative chemical complementation was performed in a different yeast strain, showing the versatility of the system, useful for performing chemical complementation with various selectable markers. This system may be extended to the ~75 human nuclear receptor proteins, plus nuclear receptors from other organisms, and the coactivators and corepressors with which they interact.

Embodiments of the present disclosure comprise chemical complementation systems focusing on one small molecule target ligand and utilize the power of genetic selection to reveal proteins within the library that bind and activate transcription in response to that small molecule. Functional receptors from a large pool of non-functional variants can be isolated, even from a non-optimized library.

Chemical complementation is a method which links survival of yeast to the presence of a small molecule. This process allows high-throughput testing of large libraries. Hundreds of thousands to billions of variants can be assayed in one experiment without the spatial resolution necessary for traditional screening methods (e.g., no need for one colony per well). Yeast can be spread on solid media and, through the power of genetic selection, cells expressing active

variants will grow into colonies. Survivors can then be spatially resolved (e.g. transferred to a microplate, one colony per well) for further characterization, decreasing the time and effort required to find new ligand-receptor pairs.

In one embodiment, among others, chemical complementation identifies nuclear receptors with a variety of responses to a specific ligand. Nuclear receptors that activate transcription in response to targeted molecules and not to endogenous compounds have several additional potential applications. The ability to switch a gene on and off in response to any desired compound can be used to build complex metabolic pathways, gene networks, and to create conditional knockouts and phenotypes in cell lines and animals. This ability can also be useful in gene therapy and in agriculture to control expression of therapeutic, pesticidal, or other genes. A variety of responses would be useful in engineering biosensor arrays: an array of receptors with differing activation profiles for a specific ligand could provide concentration measurements and increased accuracy of detection.

The ability to engineer proteins that activate transcription in response to any desired compound with a variety of activation profiles will provide a general method of identifying enzymes. Receptors that bind the product of a desired enzymatic reaction can be used to select or screen for enzymes that perform this reaction. The enzymes may be natural or engineered. The stringency of the assay can be adjusted by using ligand-receptor pairs with lower or higher EC_{50} . The lack of a general system for genetic selection is currently the limiting step for directed evolution of enzymes.

The human retinoid X receptor (RXR) is a ligand-activated transcription factor of the nuclear receptor superfamily. RXR plays an important role in morphogenesis and differentiation and serves as a dimerization partner for other nuclear receptors. Like most nuclear receptors, RXR has two structural domains: the DNA binding domain (DBD) and the ligand binding domain (LBD), which are connected by a flexible hinge region. The DBD contains two zinc modules, which bind a sequence of six bases. The LBD binds and activates transcription in response to multiple ligands including phytanic acid, docosaheptaenoic acid and 9-*cis* retinoic acid (9cRA). RXR is a modular protein; the DBD and LBD can function independently. Therefore, the LBD can be fused to other DBDs and retain function. A conformational change is induced in the LBD upon ligand

binding, which initiates recruitment of coactivators and the basal transcription machinery resulting in transcription of the target gene.

Nuclear receptors have evolved to bind, and activate transcription in response to, a variety of small molecule ligands. The known ligands for nuclear receptors are chemically diverse, including steroid and thyroid hormones, vitamin D, prostaglandins, fatty acids, leukotrienes, retinoids, antibiotics, and other xenobiotics. Evolutionarily closely related receptors (e.g., thyroid hormone receptor and retinoic acid receptor) bind different ligands, whereas some members of distant subfamilies (e.g., RXR and retinoic acid receptor) bind the same ligand. This diversity of ligand-receptor interactions demonstrates the versatility of the fold for ligand binding and suggests that it should be possible to engineer LBDs with a large range of novel specificities.

The crystal structure of RXR bound to 9cRA elucidates important hydrophobic and polar interactions in the LBD binding pocket. In one embodiment, a subset of 20 hydrophobic and polar amino acids within 4.4 Å of the bound 9cRA are varied to make a library. These residues in RXR are good candidates for creating variants that bind different ligands through site directed mutagenesis, because side chain atoms, not main chain atoms, contribute the majority of the ligand contacts. A library of RXR LBDs with all 20 amino acids at each of the 20 positions in the ligand-binding pocket screened against multiple compounds could potentially produce many new ligand-receptor pairs. However, the number of possible combinations ($20^{20} = 10^{26}$) renders saturation mutagenesis impractical for constructing a complete library.

Codon randomization creates protein libraries with mutations at specific sites. In one embodiment, a modified version of the Sauer codon randomization method to create a library of binding pocket variants of RXR is provided. This library allowed exploration of a vast quantity of sequence space in a minimal amount of time.

Chemical complementation allows testing for the activation of protein variants by specific ligands using genetic selection. In one embodiment LG335 was used, a synthetic retinoid-like compound, as a model for discovery of ligand-receptor pairs from large libraries using chemical complementation. LG335 was previously shown to selectively activate an RXR variant and not activate wild-type RXR. Combining chemical complementation with a large library of protein

variants decreases the time, effort, and resources necessary to find new ligand-receptor pairs.

Enzyme Engineering

One embodiment provides methods and compositions for engineering a polypeptide, for example an enzyme, to produce or interact with a desired molecule. Generally, a desired molecule of interest (or the reaction product) is chosen, and a target nuclear receptor is also chosen. After the target molecule and the target nuclear receptor are selected, modifications to the target nuclear receptor can be designed. For example, the X-ray structure of the target nuclear receptor can be loaded into a modeling program, including, but not limited to Insight® or Flexx®, along with the structure of the desired target molecule. Specific *in silico* interactions of the target receptor with the target molecule/ligand can be analyzed and those amino acids that may contribute the ligand binding can be noted for modification. Generally, a nuclear receptor is selected that has at least a detectable amount of interaction with the target molecule or ligand or a binding pocket of a similar size and shape. The interaction can then be modulated as desired by creating a library of modified receptors.

To create the library, site-specific codon randomization can be used. It will be appreciated that any process for generating a library of modified receptors can be used. Site-specific codon randomization involves modifying the amino acids identified through modeling as having or believed to have direct or indirect interactions with the ligand. When producing or designing the oligonucleotide, in place of those amino acids, there will be a degenerate code based on the combination of nucleotides that are desired. For example, if the modification can be a change from alanine to a cysteine, leucine, phenylalanine, isoleucine, threonine, serine, valine and methionine. The nucleotide sequence for the alanine is GCC and to possibly incorporate all of the desired amino acids mentioned above, the following changes in each position must be made:


G	C	C
1	2	3
T	T	
A		
G		G
	C	C

The oligonucleotide can be designed to have either a T, A, or G in the first position, a T or C in the second position, and a G or C in the third position. For example, if a **TTG** (one of the combinations above) is in place of the **GCC**, that
5 would incorporate a *leucine* instead of the *alanine*. Therefore, when the oligos are ordered, you would order them such that you get the possibility of a T, A, or G in the first position, a T or C in the second position, and a G or C in the third position. The oligonucleotides may be designed to include insertions or deletions. The oligonucleotides have ends that are homologous to the vector in
10 which the gene will be introduced to.

In one embodiment, to create a receptor library, the vector into which the gene will be incorporated will be cut with restriction enzymes, deleting a fragment of the wild-type gene. Oligonucleotides will be designed with homologous ends to the vector as mentioned above, but these oligonucleotides
15 will also be designed such that they overlap each other. The overlapping ends will hybridize to each other, and using for example the enzyme Klenow, the ends are filled in. Then using the polymerase chain reaction (PCR) the full gene or a fragment thereof will be amplified. After both of these products are made, these genes will be introduced into chemical complementation. The vector and gene
20 will be introduced into yeast using transformation protocols, for example protocols introduced by Gietz and co-workers. During transformation, the vector and gene or gene fragment will homologously recombine, and the various receptor mutants will be expressed.

To select for variants that bind the desired small molecule, chemical
25 complementation is be used. Chemical complementation is a general method of linking any small molecule to genetic selection. Chemical complementation is a new derivative of the yeast two-hybrid system, a three-component system that in one embodiment comprises a human nuclear receptor protein, its coactivator protein, and a small molecule ligand, where the nuclear receptor and coactivator
30 associate and activate transcription only in the presence of the ligand. An exemplary yeast strain contains a Gal4 response element fused to the ADE2 gene. If adenine is not provided in the medium, the yeast will not be able to survive unless they are able to make their own, and to do that, expression of ADE2 needs to be activated. The following exemplary plasmids can be utilized:

1st plasmid encodes a fusion protein of the Gal4 DNA binding domain (Gal4 DBD) fused to the variant receptor ligand-binding domain (LBD); the other fusion protein comprises a human coactivator protein fused to the Gal4 activation



domain. In the presence of ligand, the

25 ligand will bind to the variant receptor
ligand-binding domain and the Gal4 DNA
binding domain will bind to the Gal4

30 response element. This will cause the protein to undergo a conformational
change, and will recruit the coactivator fused to the Gal4 activation domain. This,
in turn, will result in RNA polymerase being recruited and activation of
transcription of the downstream gene.

The transformed yeast from above will be plated onto plates containing


35 the desired small molecule. Through chemical complementation, the variant
receptor that is able to bind the desired molecule and activate the *ADE2* gene
allowing that yeast colony to grow. The plasmid from that colony will be rescued
and sequenced and an engineered receptor will be identified and will be carried
on to the next step. It will be appreciated that there may be many variant

40 receptors that allow the yeast to grow without binding the targetted ligand. For
example, they may be constitutively active or bind an endogenous small
molecule. These receptors may be identified through screening without the
targetted ligand. Alternatively, they may be removed from the library by negative
genetic selection on media without the targetted ligand, either before or after

chemical complementation. Once an engineered receptor has been created, this gene can be integrated into the yeast genome, for example via homologous recombination. This will create a new strain that will be used in the following process.

Once the receptor that can bind the small molecule has been identified, individual enzymes or a library of enzymes can be evaluated to generate the product of interest. Libraries of naturally occurring enzymes, for example expression cDNA libraries, may be evaluated. Also, libraries of enzymes can be created using a number of mutagenic protocols, such as DNA shuffling, RACHITT, Error-Prone PCR, to name a few. For example, an enzyme that is suspected of interacting with the target molecule can be selected and mutagenized with conventional techniques. Alternatively, yeast or microorganisms can be randomly mutated.

In
chemical




one
embodiment,

complementation is used to identify the engineered enzyme. In this embodiment the library of engineered enzymes will be introduced into the yeast strain transformed with the modified nuclear receptor described above. This yeast strain has a variant receptor integrated into its genome, and the variant receptor is able to bind the product molecule. Once the engineered enzymes have been transformed into the yeast strain, the yeast will be spread onto selective plates

(for example plates lacking adenine) containing the reactants involved in the enzymatic reaction that can be used to synthesize the missing product. The yeast will be able to take the reactants and if the yeast express an engineered enzyme that can convert the reactants to the reaction product, then the yeast will survive. The yeast will survive because the reaction product will be able to bind to the variant receptor, and activate transcription of the *ADE2* gene or other selection gene. The DNA from the yeast colony that grew will be rescued and sequenced.

Target compounds that serve as ligands can be selected from any variety of natural or synthetic compounds. In one embodiment, natural products with agricultural or medicinal applications can be selected as target compounds. The search for natural products as potential agrochemical agents has increased due to the demand for crop protection chemicals. In 1990, the world market value of pesticides totaled nearly \$23 billion. Synthetic chemical pesticides are used to protect crops but several developments have triggered the search for alternative compounds. First, resistance has developed against synthetic chemical pesticides. Second, concern has arisen regarding potential human health risks. Third, there is a growing awareness of environmental damage, such as contamination of soil, water, and air. New environmentally friendly methods are being pursued to rectify these problems. In one embodiment of the present disclosure, the disclosed methods can be used to identify new prototype pesticides in natural products produced by microorganisms, for example, which



are perceived as more environmentally friendly and acceptable. The natural products would be applied as the synthetic chemical pesticides have been or the biosynthetic genes would be expressed in transgenic plants. This strategy has been widely applied using the *Bacillus*

thuringiensis toxin. In another embodiment, genes for toxins are delivered to target pest species using insect-specific viruses that leave beneficial insects

unharmful. These "greener" technologies require not only identification of active natural products but also the genes for their biosynthesis. With these applications in mind, and because of their availability, three compounds have been chosen as target ligands. Barbamide and jaspamide are relevant to the agricultural industry. Resveratrol has antiviral, antimicrobial, and anticancer effects.

Barbamide is a natural product from the marine cyanobacterium, *Lyngbya majuscula*. From 295 g of algae, 258 mg of pure barbamide can be isolated. This chlorinated lipopeptide has potent molluscicidal activity. The gene cluster for barbamide biosynthesis from *L. majuscula* has been cloned and analyzed. An ~26 kb region of DNA from this organism specifies the biosynthesis of barbamide. The gene cluster revealed 12 open reading frames and it is believed that barbamide is synthesized from acetate, L-phenylalanine, L-cysteine, and L-leucine. Polyketide synthase and non-ribosomal peptide synthetase modules accomplish biosynthesis. A trichloroleucine intermediate is involved, but an unresolved issue is its transfer between modules. The total synthesis of barbamide has been reported.

Jaspamide was isolated from various marine sponges and exhibits insecticidal (against *Heliothis virescens*) and fungicidal activity (against *Candida albicans*). It is completely inactive against a series of Gram negative and Gram-positive bacteria. From 700 g of sponge tissue, 80 mg of pure jaspamide was isolated. The biosynthetic pathway has not been elucidated, but its structure suggests polyketide synthase and non-ribosomal peptide synthetase modules. Since it is a fungicide, a bacterial chemical complementation system for engineering nuclear receptors and discovering the genes involved in the biosynthesis of this compound would be used.

Resveratrol is a stilbene phytoalexin that is produced in at least 72 plant species. Phytoalexins are low molecular weight antimicrobial metabolites that are produced by plants for protection against a wide range of pathogens. Some nuclear receptors are known to bind resveratrol, making the DNA shuffling approach to engineer a receptor highly relevant. This compound is commercially available on the gram scale.

Development of an amine dehydrogenase

Another embodiment provides a method for producing an enzyme, for example NAD⁺-dependent (S)-amino acid dehydrogenase.

The enzyme can preferentially produce single optical isomer products and methods and compositions for generating

between optical isomers of a compound. Variants can be achieved through

transcription of an essential gene.

Whereas the first target is a model, the second phase is an acetophenone

Conceptually, a concise and efficient method for producing products, for example amines, from

case ketones and uses ammonium formate. The method uses

yield and selectivity with concomitant recycling of NAD(P)⁺ to NAD(P)H using

enzymes such as formate dehydrogenase (FDH).

The starting enzyme is typically examined for, albeit small, levels of activity against a substrate, for example the ketone substrate in a high ammonia

environment, either i) in water/liquid ammonia-mixtures, or ii) in saturating concentrations of ammonium formate or ammonium carbonate. A sensitive

assay can be employed to check for NADH consumption such as formation of formazan ($\lambda_{\text{max}} = 450 \text{ nm}$). In this embodiment, an (S)-amino acid

dehydrogenase, either PheDH from *Rhodococcus rhodocrous* or LeuDH from

Bacillus stearothermophilus, an (R)-AmDH can be developed through change of substrate specificity. Diversity is generated within the respective gene through

both random mutagenesis and recombination. Selection via binding of the product to a nuclear receptor with subsequent transcriptional control is chosen

as the strategy to assay for successful variants.

Nuclear receptors PXR, BXR, and RAR can be used for engineering (R)-amine activated transcription with the disclosed methods and compositions. For

example, these nuclear receptors can be engineered to activate the transcription of the essential metabolic gene *ADE2* in response to the (R)-amines in the




Figure 1. Scheme for using nuclear receptors with genetic selection strategy for the directed evolution of amine dehydrogenases (AmDH). The nuclear receptor is a dimer bound to DNA at the Gal4 response element (GalRE) through the Gal4 DNA binding domain (DBD), regulating transcription of an essential gene (either *HIS3* or *ADE2*). First, a nuclear receptor ligand-binding domain (LBD) is engineered to activate transcription in response to the desired (R)-amine. Second, libraries of AADH are transformed into the microbe and grown on media supplemented with the appropriate ketone. Only microbes with a functional AmDH that converts the ketone into the (R)-amine survive.

modified *Saccharomyces cerevisiae* strain PJ69. PXR is chosen because of its broad substrate specificity. BXR is chosen because it is already known to activate transcription in response to amines. Random and structure-based approaches of creating libraries to engineer the nuclear receptors for (R)-amine
5 activated growth through genetic selection can be used. Receptors for multiple (R)-amines will be engineered in parallel by selecting each library on multiple selective plates with the appropriate (R)-amine. Optionally, negative selection to genetically select libraries against enzymes that make an S-enantiomer product then select for the production of the R-enantiomer (or vice-versa) can be used. A
10 nuclear receptor library for the (R)-amine ligand can be synthesized. Additionally, the (R)-amine ligand can be synthesized *in vivo* by an expressed AmDH from the ketone precursor supplemented within the growth medium. A mutant PheDH library can then be screened for *in vivo* synthesis of (R)-amines. In this overall scheme, the power of genetic selection is used to detect
15 biocatalytic synthesis of amines. Utilizing genetic selection means that each member of the library does not need to be screened, only functional AmDH appear because they allow the microbe to grow and form a colony. Furthermore, catalysis is *directly selected*, as opposed to some related but indirect property (like transition state binding). Genetic selection coupled with the broad ligand
20 specificity of nuclear receptors creates a process to rapidly improve biocatalysts for more efficient synthesis of enantiomerically pure compounds.

Selected transformants can be optimized through successive rounds of directed evolution. Further mutant libraries of PheDH/LeuDH enzymes can be screened for *in vivo* synthesis of (R)-amine. Mutant AmDH enzymes can be
25 expressed and further studied for shifts in substrate specificity and changes in kinetic reaction rates.

Fig. 10 depicts another embodiment for the identification of selective receptor modulators (analogous to selective estrogen modulators). In this embodiment, the human nuclear receptor coactivator ACTR is fused to the Gal4
30 activation domain (ACTR:GAD). Additionally, the human nuclear receptor coactivator SRC1 is fused to a yeast repression domain (SRC1:RD). In the presence of an agonist, these coactivator fusion proteins compete for expression of the *HIS3* gene. The *HIS3* gene encodes imidazoleglycerolphosphate dehydratase. In the presence of an agonist that recruits both coactivators

equally, the yeast probably will produce enough histidine to survive. Adding the inhibitor 3-AT to the plates raises the threshold of enzyme that must be produced to permit growth. Compounds that selectively favor the RXR-ACTR interaction over the RXR-SRC-1 interaction will allow yeast to grow.

5 **Fig. 11** is a diagram of another embodiment incorporating negative chemical selection. Human nuclear receptor coactivator, ACTR is fused to the Gal4 activation domain (ACTR:GAD). The Gal4 DBD is fused to the nuclear receptor LBD (GBD:RXR). The Gal4 DBD binds to the Gal4 response element, regulating transcription to the *URA3* gene. The *URA3* gene codes for orotidine-
10 5'-phosphate decarboxylase, an enzyme in the uracil biosynthetic pathway. This gene can be used for both positive and negative selection. For positive selection, yeast expressing this gene will survive in the absence of uracil in the media. For negative selection, 5-fluoroorotic acid (FOA) is added to the media. Expression of orotidine-5'-phosphate decarboxylase converts FOA to the toxin 5'-
15 fluorouracil, which kills the yeast. Libraries of small molecules can be screened in a high-throughput assay in wells containing an agonist and FOA. Antagonists will allow yeast to grow.

Fig. 12 is a diagram illustrating still another embodiment comprising isotype specific nuclear receptor agonists are. Each isotype can be fused to a
20 different DBD controlling expression of different genes. The isotype for which an agonist is sought is fused to the Gal4 DBD to control expression of *ADE2* (for positive chemical complementation). The isotype against which selectivity is desired, is fused to the GCN4 DBD to control expression of the *URA3* gene (for negative chemical complementation). Libraries of small molecules are screened
25 in individual wells of a 384-well plate. Compounds that do not activate the receptor will not allow the yeast to grow. Compounds that agonize both isotypes will kill the yeast. Only compounds that agonize RXR α , and either do not bind or antagonize RXR β will allow yeast to grow.

Fig. 13 shows another embodiment in which a human nuclear receptor
30 coactivator, ACTR, is fused to the Gal4 activation domain (ACTR:GAD). The Gal4 DBD is fused to the nuclear receptor LBD (GBD:RXR). The Gal4 DBD binds to the Gal4 response element, regulating transcription of the *ADE2* gene. Upon binding of the ligand, the LBD of the nuclear receptor undergoes a conformational change, which recruits the ACTR:GAD fusion protein. This

brings the Gal4 AD and Gal4 DBD into close proximity activating transcription of the *ADE2* gene. For clarity only one ACTR:GAD protein is shown binding one GBD:RXR. Libraries of small molecules are screened in individual wells of a 384-well plate. Agonists will allow yeast to grow.

5

Materials and Methods

Ligands. 9-cis retinoic acid (MW=304.44 g/mol) was purchased from ICN Biomedicals.

LG335 Synthesis

10 **3-(1-Carbonyl)propyl-5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthylene**
2,5-dimethyl-2,5-hexanediol (5.0 g, 34 mmol) was dissolved in anhydrous benzene (150 mL). AlCl_3 (5.0 g, 38 mmol) was added slowly while the mixture was stirred in an ice bath, followed by stirring at room temperature for 1 hour. Another portion of AlCl_3 (5.0 g, 38 mmol) was then added and the reaction was
15 heated to 50 °C and stirred overnight. The brown solution was poured over iced 0.4 M HCl (50 mL) and extracted with ether (3 x 50 mL). The organic layer was then sequentially washed with water, saturated aqueous NaHCO_3 , and brine (80 mL each) and dried (MgSO_4). The solvent was removed in vacuo to afford 6.2 g of a yellow liquid (2).

20 The crude product was then mixed with propionyl chloride (3.2 mL, 37 mmol) and the resulting solution added dropwise to a mixture of AlCl_3 (5.0g, 38 mmol) in dichloroethane (20mL) while maintaining the temperature between 20 and 25 °C. The mixture was stirred for 2 hours at room temperature, at which point it was quenched by pouring carefully over ice. The reaction mixture was
25 then extracted methylene chloride (3 x 10 mL). The organics layers were then combined, washed with water and saturated aqueous NaHCO_3 the volatiles removed by rotary evaporation. The product was purified by silica gel column chromatography eluting with hexanes:chloroform (4:1, then 1:1) to yield 6.9 g (28 mmol, 73%) of product as a yellow oil (3, 4).

30 **3-Propyl-5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthylene**
3-(1-Carbonyl)propyl-5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthylene (1.0 g, 4.1 mmol) in MeOH (10 mL), H_2O (1 mL), and conc. HCl (3 drops) was treated with 10% Pd/C (144 mg) and subjected to catalytic hydrogenation conditions at 60 psi while heating gently overnight. When the reaction was considered complete (Rf

= 0.76, 5% EtOAc in hexanes) it was filtered through a celite pad and rinsed with MeOH (10 mL) and hexane (50 mL). Water (1 mL) was then added to the filtrate and the organic phase separated and washed with brine (2 x 20 mL). The aqueous layer was washed with hexanes (2 x 20 mL). The organic layers were dried (Na₂SO₄), filtered and the volatiles removed by rotary evaporation to produce 510 mg (2.2 mmol, 54%) of a colorless oil (5).

4-[(3-Propyl-5,5,8,8-tetramethyl –

5,6,7,8-tetrahydro-2-naphthyl)carbonyl]benzoic Acid (LG335)

3-Propyl-5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthylene (2.2 g, 9.5 mmol) and chloromethyl terephthalate (2.0g, 10 mmol) were dissolved in dichloroethane (20 mL) and FeCl₃ (80 mg, 490 μmol) was added. The reaction mixture was stirred at 75 °C for 24 hours. The reaction was then cooled and MeOH (20 mL) added. The resulting slurry stirred for 7 hours at room temperature, filtered and rinsed with cold MeOH (20 mL) to result in 2.1 g (5.5 mmol, 58%) of white crystals (6).

The crystals (107 mg, 280 μmol) were stirred in MeOH (2 mL), to which 5N KOH (0.5 mL) was added. This mixture was refluxed for 30 minutes, cooled to room temperature and acidified with 20% aqueous HCl (0.5 mL). The MeOH was evaporated and the residue was extracted with EtOAc (2 x 5 mL). The organic layers were combined and dried (MgSO₄) and filtered. The filtrate was treated with hexane (10 mL) and reduced in volume to 2 mL. After standing overnight the resulting crystals were collected to provide 39 mg (103 μmol, 37%) as a white powder (1). mp 250-252 °C; ¹H NMR (CDCl₃) δ 0.88 (t, 3H, -CH₂CH₂CH₃), 1.20 (s, 6H, CH₃), 1.32 (s, 6H, CH₃), 1.55 (dt, 2H, -CH₂CH₂CH₃), 1.69 (s, 4H, CH₂), 2.65 (t, 2H, -CH₂CH₂CH₃), 7.20 (s, 1H, Ar-CH) 7.23 (s, 1H, Ar-CH), 7.89 (d, 2H, Ar-CH), 8.18 (d, 2H, Ar-CH); MS (EI POS) m/z mass for C₂₅H₃₀O₃: Calc. 378.2189, Found 378.2195; Anal. for C₂₅H₃₀O₃: Calc. C:79.33, H:7.99, Found C:79.10, H:7.96.

Expression Plasmids. pGAD10BAACTR, pGBT9Gal4, pGBDRXRα, pCMX-hRXR, and pCMX-βGAL have been described. pCMX-hRXR mutants were cloned from pGBDRXR vectors using *Sall* and *PstI* restriction enzymes and ligated into similarly cut pCMX-hRXR vectors. pLuc_CRBPII_MCS was constructed as below. All plasmids have been confirmed through sequencing.

pGBDRXR α was cut with *Sma*I and *Nco*I, filled in, and blunt-end ligated to eliminate 153 amino acids of the RXR DBD. A *Hind*III site in the tryptophan selectable marker was silently deleted and the sole remaining *Hind*III site was cut, filled in, and blunt-end ligated to remove the restriction site. Unique *Hind*III and *Sac*I sites were inserted into the RXR LBD gene and *Mfe*I and *Eco*RI sites were removed from the plasmid using QuikChange Site-Directed Mutagenesis (Stratagene, La Jolla, CA) to create pGBDRXR α L-SH-ME.

pLuc_CRBPII_MCS was made by site-directed mutagenesis from pLucMCS (Stratagene, USA). Site-directed primers were designed to incorporate a CRBPII response element in the multiple cloning site (MCS), controlling transcription of the firefly luciferase gene.

Plasmids expressing the fusion protein of the Gal4 activation domain with the coactivators are based on the commercial plasmid pGAD10 (Clontech, USA). The pGAD10 vector contains the Gal4 activation domain (residues 491-829) fused to a multiple cloning site (MCS) and uses a leucine marker. Additional restriction enzyme sites were added to the MCS of the plasmid via site directed mutagenesis. Primers were designed to add the following restriction enzymes: *Nde*I, *Eag*I, *Ecl*XI, *Not*I, *Xma*III, *Xma*I, and *Sma*I, forming a new plasmid known as pGAD10BA. (Figure 17) This plasmid was sequenced and used for specific interaction studies mentioned in the results.

pCMX-ACTR, the expression plasmid for the human nuclear receptor coactivator ACTR, was a kind gift from Dr. Ron Evans (Salk Institute for Biological Studies, La Jolla, CA). pCR3.1hSRC-1, the expression plasmid for the human nuclear receptor coactivator SRC-1, was a kind gift from Dr. Bert O'Malley (Baylor College of Medicine, Houston, TX). Both ACTR (residues 1-1413) and SRC-1 (residues 54-1442) genes were amplified via PCR with primers that contained *Bgl*II and *Not*I sites. The PCR products were digested with the two restriction enzymes and cleaned using the Zymo "DNA Clean and Concentrator Kit" (Zymo Research, Orange, CA) spin columns, pGAD10BA was digested with *Bgl*II and *Not*I and ligated with both the ACTR and SRC-1 products. Ligations were transformed into Z-competent (Zymo Research, Orange, CA) XL 1-Blue cells (Stratagene, La Jolla, CA).

Transformants were rescued and sequenced. The final plasmids are called pGAD10BAACTR and pGAD10BASRC1.

Plasmid Construction. The zero background plasmid, pGBDRXR:3Stop, was constructed using QuikChange Site-Directed Mutagenesis with pGBDRXR α L-SH-ME as the template and the 3Stop insert cassette (described below) as primers.

The 3Stop insert cassette was synthesized using PCR from eight oligonucleotides (Fig. 16). All PCRs were done using 2.5 U Pfu Polymerase (Stratagene, LaJolla, CA), 1x Pfu buffer, 0.8 mM dNTPs, 50 ng of pGBDRXR α L-SH-ME as a template, 125 ng of primers and sterile water to make 50 μ L. First, four small cassettes were synthesized in reactions containing the following primers: Cassette 1, F (5'-CGGAATTTC CATGGGC-3') (SEQ ID NO. 1), BPf (5'-CTCGCCGAAC GACCCGGTCA CCGCATGCCA CTAGTGG-3') (SEQ ID NO. 2), and BPr (5'-CCGCTTGGCC CACTCCACTA GTGGCATGCG GTGACC-3') (SEQ ID NO. 3); Cassette 2, BPf, BPr, SEf (5'-CGGGCAGGCT GGAATGAGCT CCTCGACGGA ATTCTCC-3') (SEQ ID NO. 4), and SEr (5'-CAGCCCGGTG GCCAGGAGAA TTCCGTCGAG GAGCTC-3') (SEQ ID NO. 5); Cassette 3, SEf, SEr, AMf (5'-CTCTGCGCTC CATCGGGCTT AAGTGCCAC CAATTGACAC-3') (SEQ ID NO. 6), and AMr (5'-CTCCAGCATC TCCATAAGGA AGGTGTCAAT TGGTGGGCAC TTAAGC-3') (SEQ ID NO. 7); Cassette 4, AMf, AMr, and R (5'-CAAAGGATGG GCCGCAG-3') (SEQ ID NO. 8). The cassettes were cleaned with either the DNA Clean and Concentrator-5 (Zymo Research, Orange, CA) or the Zymoclean Gel DNA Recovery Kit (Zymo Research, Orange, CA) depending on product purity. The four cassettes were used to make the final 3Stop insert cassette in a PCR that contained each cassette, primers F and R, dNTPs, Pfu Polymerase, and sterile water to a final volume of 50 μ L. The 3Stop cassette was cleaned using the Zymoclean Gel DNA Recovery Kit.

Insert Cassette Library Construction. The library of insert cassettes with randomized codons was constructed in a similar manner as above. The four

cassettes (FBP, BPSE, SEAM and AMR) were made in the following ways (Supporting Information Fig. 7b).

For the FBP cassette, oligos BP1 (5'-GGCAAACATG GGGCTGAACC CCAGCTCGCC GAACGACCCG GTCACC-3') (SEQ ID NO. 9), BP2 (5'-GCCCACTCCA CTAGTGTGAA AAGCTGTTTG TC (A, C, or T)(A or G)(C or G)(A, C, or T)(A or G)(C or G)TT GGCA(A, C, or T)(A or G)(C or G)GTT GGTGACCGGG TCGTTCG-3') (SEQ ID NO. 10), BP3 (5'-CTTTTCACAC TAGTGGAGTG GGCCAAGCGG ATCCCACACT TCTCAGAG-3') (SEQ ID NO. 11), and BP4 (5'-GGGGCAGCTC TGAGAAGTGT GGGATCCG-3') (SEQ ID NO. 12) were mixed with TE containing 100 mM NaCl to bring the total volume to 50 μ L. The mixture was heated to 95 °C for 1 minute, then slowly cooled to 10 °C. The annealed mixture was combined with EcoPol Buffer, dNTPs, ATP, Klenow (NEB, Beverly, MA), T4 DNA ligase (NEB, Beverly, MA) and sterile water to 200 μ L, and kept at 25°C for 45 min before heat inactivation at 75°C for 20 minutes. The product was cleaned with DNA Clean and Concentrator-5 to make the BP cassette. Next, BP cassette was combined with Pfu Buffer, pGBDRXR:3Stop, oligo F, dNTPs, Pfu polymerase, and sterile water to make 50 μ L for a PCR. The final FBP product (300bp) was purified using the Zymoclean Gel DNA Recovery Kit.

BPSE was made in two consecutive PCRs. First, SE1 (5'-GCAGGCTGGA ATGAGCTCCT C(A, G, or T)(C or T)(G or C)GCCTCC (A, G, or T)(C or T)(G or C)TCCCACC GCTCCATC-3') (SEQ ID NO. 13) and SE2 (5'-CCGGTGGCCA GGAGAATTCC GTCCTTCACG GCGATGGAGC GGTGGG-3') (SEQ ID NO. 14) were combined with Pfu buffer, dNTPs, Pfu polymerase, and sterile water to make 50 μ L. After 5 PCR cycles, pGBDRXR:3Stop and BP were added to the reaction and the PCR was continued for 30 cycles. The product (240 bp) was purified using the Zymoclean Gel DNA Recovery Kit.

SEAM was constructed in a similar way to BPSE. SE1 and SE2 were mixed with Pfu Buffer, dNTPs, Pfu polymerase, and sterile water to 25 μ L. Simultaneously, AM1 (5'-GGCTCTGCGC TCCATCGGGC TTAAGTGCCT GGAACAT(A, G, or T)(C or T)(G or C) TTSCTTCTTC AAGCTCATCG GGG-3')(SEQ ID NO. 15) and AM2 (5'-GCATCTCAAT AAGGAAGGTG

TCAATTGTGT GTCCCCGATG AGCTTGAAGA A-3') (SEQ ID NO. 16) were combined with Pfu Buffer, dNTPs, Pfu polymerase, and sterile water to 25 μ L. After 5 cycles, these two reactions were mixed and pGBDRXR:3Stop was added. The PCR was continued for 30 cycles. The PCR product (460 bp) was purified using the Zymoclean Gel DNA Recovery Kit.

The AMR cassette was made similarly to FBP. AM1 and AM2 were mixed with TE containing 100 mM NaCl to make 50 μ L, heated to 95°C for 1 minute, then slowly cooled to 10°C. The annealed mixture was combined with EcoPol Buffer, dNTPs, Klenow, and sterile water to 200 μ L, and kept at 25°C for 45 min before heat inactivation at 75°C for 20 minutes. The product (AM) was precipitated with isopropanol. Next, AM and R were combined with Pfu buffer, pGBDRXR:3Stop, dNTPs, Pfu Polymerase, and sterile water to make 50 μ L for a PCR. The product (140 bp) was purified using the Zymoclean Gel DNA Recovery Kit.

The four cassettes (FBP, BPSE, SEAM, and AMR) were combined in a PCR to make the library of randomized insert cassettes (6mutIC). The library was cleaned using Bio-Spin 30 columns (Bio-Rad Laboratories, Hercules, CA).

Yeast selection plates and transformation. Synthetic complete (SC) media and plates were made as previously described (7). Selective plates were made without tryptophan (-Trp) and leucine (-Leu) or without adenine (-Ade), tryptophan (-Trp) and leucine (-Leu). Ligands were added to the media after cooling to 50 °C.

The randomized cassette library was homologously recombined into the pGBDRXR:3Stop plasmid using the following method. pGBDRXR:3Stop was first digested with *BssHII* and *EagI* (NEB, Beverly, MA), and then treated with calf intestinal phosphatase (NEB, Beverly, MA), to make a vector cassette. Vector cassette (1 μ g) and 6mutIC (9 μ g) were transformed according to Geitz's transformation protocol (8) on a 10X scale into the PJ69-4A yeast strain, which had previously been transformed with a plasmid (pGAD10BAACTR) (manuscript submitted) expressing the nuclear receptor coactivator ACTR fused to the yeast Gal4 activation domain. Homologous regions between the vector cassette and the insert cassette allow the yeast to homologously recombine the insert

cassette with the vector cassette forming a circular plasmid with a complete RXR LBD gene. The transformation mixture (1 mL) was spread on each of 10 large plates of SC -Ade -Trp -Leu media containing 10 μ M LG335. The transformation mixture (2 and 20 μ L) was also spread on SC -Trp -Leu media. These plates
 5 were grown for 4 days at 30 °C.

Molecular Modeling. Docking of LG335 in to modified binding pockets was done using the InsightII module Affinity. The wild type RXR with 9cRA crystal structure (9) was modified using the Biopolymer module residue replace tool to
 10 make mutations in the binding pocket that corresponded to the mutations in variants I268;I30A;F313A;L436F, I268V;A272V;I310L;F313M, and I268A;I310S;F313A;L436F. The ligand was placed in the binding pocket by superimposing the carboxylate carbon and two carbons in the tetrahydronaphthalene ring of LG335 onto corresponding carbons of 9cRA in the
 15 crystal structure. A Monte Carlo simulation was performed first, followed by Simulated Annealing of the best docked conformations.

Library Evaluation

To evaluate the efficiency of library creation and selection we take a binary
 20 approach— either the sequence is or is not a designed sequence. Eq. 1 is the relevant binomial distribution for statistical evaluation of the libraries.

$$P = \frac{(N-1)!}{(k-1)!(N-k)!} p^k (1-p)^{N-k} \quad (1)$$

In Eq. 1 N is the number of sequenced plasmids; k is the number of background or designed plasmids; p is the frequency of the occurrence of either background or designed plasmid; and P is the measure of certainty. Applying Eq. 1 to the
 25 libraries, we conclude with 95% certainty that the unselected library is at least 72% background and the selected library is at least 78% designed sequences.

Genotype Determination. Plasmids were rescued using either the Powers method (www.fhcrc.org/labs/gottschling/yeast/yplas.html) or the Zymoprep Kit
 30 (Zymo Research, Orange, CA). The plasmids were then transformed into Z-competent (Zymo Research, Orange, CA) XL1-Blue cells (Stratagene, La Jolla,

CA). The QIAprep Spin Miniprep Kit (Qiagen, Valencia, CA) was used to purify the DNA from the transformants. These plasmids were sequenced.

Quantitation Assays

- 5 **Solid Media.** The rescued plasmids were transformed into PJ69-4A containing the pGAD10BAACTR plasmid and plated on (SC) -Trp -Leu media. These plates were grown for 2 days at 30 °C.

Colonies were streaked onto the following media: SC, SC -Trp -Leu, SC -Ade -Trp -Leu, SC -Ade -Trp -Leu plus increasing concentration of LG335 or
10 9cRA from 1 nM to 10 µM.

Liquid Media. The method used for quantitation was modified from a method developed by Miller and known in the art.

- 15 **Mammalian Luciferase Assay.** Performed with HEK 293 cells as previously described, and known in the art.

Streaking cells onto adenine selective plates using PJ69-4A.

- Yeast transformants containing the plasmids were streaked onto the
20 selective plates (SC -Ade) with different ligand concentrations using sterile toothpicks. Plates were divided into sectors for the samples and controls; the control sectors contain pGBDMT and pGBT9Gal4. The same colony was used for streaking on all the plates, ending with a SC plate to confirm efficient transfer of the cells to each plate. Both selective and non-selective plates
25 were incubated at 30 °C for two days. Each set of genetic selection plates was replicated at least once.

Streaking cells onto FOA plates using MaVW3

- Yeast transformants containing the plasmids were streaked onto
30 selective plates, SC -Leu-Trp, containing 5-fluororotic acid, FOA, and different ligand concentrations. Plates were also divided into sectors, with pGBT9Gal4 and pGBDMT as controls. The same procedure was used for

streaking as for the adenine selection plates. Plates were incubated for two days. Each set of the genetic selection plates was replicated at least once.

EXAMPLES

Example 1: Library Design

5 The binding pocket of the RXR LBD is composed of primarily hydrophobic side chains plus several positively charged residues that stabilize the negatively charged carboxylate group of 9cRA. The target ligand, LG335, contains an analogous carboxylate group, so the positively charged residues were left unchanged. We hypothesized that binding affinity arises from hydrophobic
10 contacts and that specificity arises from binding pocket size, shape, hydrogen bonding, and electrostatics. The randomized amino acids were chosen based on their proximity to the bound 9cRA as observed in the crystal structure and the results of site directed mutagenesis (supporting information Fig. 14). The electrostatic interactions were held constant while the size, shape, and potential
15 hydrogen bonding interactions were varied to find optimum contacts for LG335 binding. A library of RXRs with mutations at six positions was created. At three of the positions (I268, A271, and A272) are four possible amino acids (L, V, A, and P) and at the other three positions (I310, F313, and L436) there are eight possible amino acids (L, I, V, F, M, S, A, and T). The combination of six positions
20 and number of encoded amino acids allowed testing of the library construction while keeping the library size (32,768 amino acid combinations and ~3 million codon combinations) within reasonable limits. Proline was included in the library as a negative control. Residues 268, 271, and 272 are in the middle of helix 3, which would be disrupted by the inclusion of proline. Therefore, proline residues
25 should appear at these positions only in unselected variants and not in the variants that activate in response to ligand. The substitutions at positions 268, 271, and 272 were restricted to small amino acids allowing access to the positively charged residues at this end of the pocket.

 To eliminate contamination of the library with unmutated, wild-type RXR
30 the gene was modified to create a non-functional gene, RXR:3Stop. Forty base pairs were deleted at three separate sites producing three stop codons in the coding region to create this nonfunctional gene. The deletions correspond to regions in the RXR gene where randomized codons are designed. This plasmid,

pGBDRXR:3Stop, was cotransformed into yeast with the library of insert cassettes containing full-length RXR LBD genes with randomized codons at positions 268, 271, 272, 310, 313, and 436. The insert cassettes and the plasmid contain homologous regions enabling the yeast to homologously recombine the cassette into the plasmid. Recombination repairs the deletions in the RXR:3Stop gene to make full-length genes with mutations at the six specific sites.

Example 2: Library selection.

To limit the number of variants to be screened, the library was subjected to chemical complementation (Fig. 1). Chemical complementation exploits the power of genetic selection to make the survival of yeast dependent on the presence of a small molecule. The PJ69-4A strain of *S. cerevisiae* has been engineered for use in yeast two-hybrid genetic selection and screening assays. For selection, PJ69-4A contains the ADE2 gene under the control of a Gal4 response element. Plasmids created through homologous recombination in PJ69-4A express the Gal4 DBD fused with a variant RXR LBD (GBD:RXR). A plasmid expressing ACTR, a nuclear receptor coactivator, fused with the Gal4 activation domain (ACTR:GAD), was also transformed into PJ69-4A. If a ligand causes a variant RXR LBD to associate with ACTR, transcription of the ADE2 gene is activated. Expression of ADE2 permits adenine biosynthesis and therefore, yeast survival on media lacking adenine.

A small amount of the yeast library was plated onto media (SC -Leu -Trp) selecting only for the presence of the plasmids pGAD10BAACTR (expressing ACTR:GAD and containing a leucine selective marker) and mutant pGBDRXR (expressing variant GBD:RXR and containing a tryptophan selective marker). The majority of the yeast cells transformed with the RXR library were plated directly onto SC -Leu -Trp -Ade media containing 10 μ M LG335, selecting for adenine production in response to the compound LG335. The transformation efficiency of this library into yeast strain PJ69-4A was 3.8×10^4 colonies per μ g DNA. This number includes both the efficiency of transforming the DNA into the cells and the homologous recombination efficiency. Of the approximately 380,000 transformants, approximately 300 grew on SC -Ade -Trp -Leu + 10 μ M LG335 selective media.

Example 3: Library Characterization.

Twenty-one plasmids were rescued from yeast colonies: nine from non-selective plates (SC -Trp -Leu) and twelve from selective plates (SC -Ade -Trp -Leu + 10 μ M LG335). The relevant portion of plasmid DNA from these colonies was sequenced to determine the genotype (Table 1). All nine of the plasmid sequences from the non-selective plates contained at least one deletion and are non-functional genes. Of the twelve plasmids that grew on the selective media, all contain full-length RXR LBDs with designed mutations. With 95% certainty, we conclude that the unselected library is at least 72% background and the selected library is at least 78% designed sequences (supporting information).

Table 1. Genotypes of mutants from unselected and selected libraries

Mutant	I268	A271	A272	I310	F313	L436
Unselected library						
1	Deleted	Deleted	Deleted	Deleted	Deleted	Deleted
2	Deleted	Deleted	Deleted	Deleted	Deleted	Deleted
3	GTA(V)	CCT(P)	CCT(P)	TCG(S)	TCG(S)	Deleted
4	Deleted	Deleted	Deleted	Deleted	Deleted	Deleted
5	Deleted	Deleted	Deleted	Deleted	Deleted	GCG(A)
6	Deleted	Deleted	Deleted	Deleted	Deleted	Deleted
7	Deleted	Deleted	Deleted	Deleted	Deleted	Deleted
8	Deleted	Deleted	Deleted	Deleted	Deleted	Deleted
9	Deleted	Deleted	Deleted	Deleted	Deleted	TTC(F)
Selected library						
1	GTG(V)	wtRXR	GCA	TTG(L)	ATG(M)	TTG
2	GTG(V)	wtRXR	GCA	GTG(V)	TCC(S)	TTG
3	CTA(L)	GCT	GCA	ATG(M)	GTG(V)	TTG
4	GCG(A)	wtRXR	GCA	TCC(S)	GTG(V)	TTC(F)
5	GCT(A)	GCT	GCA	GCC(A)	GCG(A)	TTC(F)
6	GCT(A)	GCT	GTT(V)	GCC(A)	GCG(A)	TTC(F)
7	CTT(L)	GCT	GCT	GTC(V)	ATC(I)	TTG
8	CTG(L)	GTG(V)	GCG	TTG(L)	TTG(L)	TTG
9	GTG(V)	GTG(V)	GCG	TTG(L)	GTG(V)	TTG
10	GTA(V)	wtRXR	GTG(V)	ATG(M)	TCC(S)	ATG(M)
11	GCG(A)	GCG	GCA	ATG(M)	GCG(A)	ACG(T)
12	GCG(A)	GCT	GCG	TCG(S)	GTC(A)	TTC(F)

Sequences condons are followed by the encoded amino acid in parentheses.

"wtRXR" indicates that the sequence corresponds to the wild-type RXR condon.

"Deleted" indicates the presence of an unmutated 35top deletion background cassette.

Example 4: Variant Characterization in Yeast.

The twelve plasmids rescued from the selective plates were retransformed into PJ69-4A to confirm that their phenotype is plasmid linked. The strain PJ69-4A was engineered to contain a Gal4 response element controlling expression of the *LacZ* gene, in addition to the *ADE2* gene. Both selection and screening were used to determine the activation level of each variant by 9cRA and LG335. The selection assay quantifies yeast growth occurring through transcriptional activation of the *ADE2* gene, while the screen quantifies β -galactosidase activity occurring through transcriptional activation of the *LacZ* gene. Although the selection assay (Fig. 2) is ~10-fold more sensitive than the screen (Fig. 3), it does not quantify activation level (efficacy) as well as the screen. In the selection assay, there is either growth or no growth, whereas the screen more accurately quantifies different activation levels at various concentration of ligand (Figs. 2 and 3). The differences will be more fully discussed in a future publication.

Three plasmids were used as controls in the screen and selection assays. The plasmids pGBDRXR α and pGBT9Gal4 were used as positive controls to which the activation level of the variants can be compared. pGBDRXR α expresses the gene for the "wild-type" GBD:RXR, which grows and is activated by 9cRA but not by LG335. pGBT9Gal4 expresses the gene for the ligand-independent yeast transcription factor Gal4 (25), which is constitutively active in the presence or absence of either ligand. The plasmid pGBDRXR:3Stop serves as a negative control. pGBDRXR:3Stop carries a non-functional RXR LBD gene; therefore, yeast transformed with this plasmid does not grow in the selection assay nor show activity in the screen. This plasmid provides a measure of background noise in both the selection and screen assays.

Both the selection and screen assays show that ten of the twelve variants are selectively activated by LG335. Results of these assays are shown in figures 2 and 3. Table 2 summarizes the transcriptional activation profiles of all twelve variants in response to both 9cRA and LG335 compared to wild-type RXR.

Table 2. EC₅₀ and efficacy in yeast and HEK 293 cells for RXR variants

Variant	9CRA				LG335			
	Yeast		HEK 293		Yeast		HEK 293	
	EC ₅₀	Eff	EC ₅₀	Eff	EC ₅₀	Eff	EC ₅₀	Eff
WT	500	100	220	100	>10,000	10	300	10
I268A;I310A;F313A;L436F	>10,000	0	>10,000	0	220	70	30	50
I268V;A272V;I310L;F313M	>10,000	10	1,600	30	40	60	1	30
I268A;I310S;F313V;L436F	>10,000	10	-	-	470	60	-	-
I268A;I310S;F313V;L436F	>10,000	0	>10,000	0	430	50	690	20
I268V;A272V;I310M;F313S;L436M	>10,000	10	>10,000	0	680	30	180	30
I268A;A272V;I310A;F313A;L436F	>10,000	0	-	-	530	30	1	-
I268L;A271V;I310L;F313L	>10,000	0	-	-	530	20	1	-
I268A;I310M;F313A;L436T	>10,000	0	>10,000	0	610	10	140	20
I268V;A271V;I310L;F313V	>10,000	0	-	-	650	10	-	-
I268L;I310V;F313I	>10,000	0	-	-	>2000	10	-	-
I268L;I310M;F313V	>10,000	20	-	-	610	20	-	-
I268V;I310V;F313S	>10,000	0	-	-	440	10	-	-

EC₅₀ values (given in nm) represent the averages of two screen experiments in quadruplicate for yeast and in triplicate for HEK 293. Efficacy (Eff; given as a percent) is the maximum increase in activation relative to the increase in activation of wild type with 10 μ M 9cRA. Values represent the averages of two screen experiments in quadruplicate for yeast and in triplicate in HEK 293.

Five variants were chosen for testing in mammalian cell culture for comparison of the activation profiles (I268A;I310A;F313A;L436F, I268V;A272V;I310L;F313M, I268A;I310S;F313A;L436F, I268V;A272V;I310M;F313S;L436M, and I268A;I310M;F313A;L436T). The genes for these variants were removed from yeast expression plasmids and ligated into mammalian expression plasmids.

Although I268L;I310M;F313V is constitutively active in the selection assay (Fig. 2n) and has high basal activity in the screen assay, both 9cRA and LG335 increase activity at micromolar concentrations (Fig. 3n). This variant may be in an intermediate conformation, with weakly activated transcription that can be improved by ligand binding. The high basal activation could also be due to a change in the conformation equilibrium with a shift towards the active conformation when ligand is not present.

I268V;I310V;F313S is constitutively active on solid media (data not shown), but shows no activation in the screen (0% Eff., Table 2, Fig. 3o) and only grows in the liquid media selection after two days (Fig. 2o). The basal activation level may be below the threshold of detection for the liquid media assays. However, it is also possible that agar, which is not present in the liquid

Activation levels and EC_{50} s correlate in yeast and HEK 293 cells (Fig. 4 and Table 2). For the majority of the variants 9cRA shows little or no activation in yeast or mammalian cells. Variant I268V;A272V;I310L;F313M is activated slightly by 9cRA in yeast, but in mammalian cells is activated to the same level as with both 9cRA and LG335 (Figs. 2, 3 and 4). With one exception, all variants tested have EC_{50} s within 10-fold in yeast and mammalian cells. However, the EC_{50} s in mammalian cells are generally lower than in yeast. We speculate that this shift is due to increased penetration of LG335 into mammalian cells versus yeast.

Subtle differences in binding pocket shape can have a drastic effect on specificity. For example, the I268V;A272V;I310L;F313M variant is activated to high levels by LG335 (60% Eff. Table 2), and is only slightly activated by 10 μ M 9cRA in yeast (Fig. 3e), yet the amino acid changes are extremely conservative. The volume difference between phenylalanine and methionine side chains is only $\sim 4 \text{ \AA}^3$ and their polarity difference is minimal (hydration potentials of the methionine and phenylalanine side chains are $-0.76 \text{ kcal mol}^{-1}$ and $-1.48 \text{ kcal mol}^{-1}$, respectively). The other mutations redistribute methyl groups within the binding pocket, with a net difference of one methyl group ($\sim 18 \text{ \AA}^3$).

The LG335-I268V;A272V;I310L;F313M ligand receptor pair also represents a 25-fold improvement in EC_{50} over the previous best LG335 receptor, Q275C;I310M;F313I (40 nM vs. 1 μ M in yeast). The Q275C;I310M;F313I variant was created using site directed mutagenesis. Subtle changes in the I268V;A272V;I310L;F313M variant produced a better ligand receptor pair than the Q275C;I310M;F313I variant. This conclusion is consistent with the observation that nuclear receptors bind ligands through an induced-fit mechanism. With current knowledge about protein-ligand interactions it is not possible to rationally design ligand-receptor pairs with specific activation profiles.

Libraries and chemical complementation are a new way to circumvent this problem and obtain functional variants with a variety of activation profiles.

Molecular modeling was used to generate hypotheses about the structural basis of ligand specificity for the variants discovered in the library. First, mutations to smaller or more flexible side chains at positions 310, and 313 are essential to provide space for the propyl group of LG335. All variants activated by LG335 have mutations at these two positions. Second, mutations to amino acids with larger side chains at position 436 sterically clash with the methyl group at the 9 position of 9cRA. This interaction may prevent helix 12 from closing properly and therefore prevent activation by 9cRA. The only variant significantly activated by 9cRA (I268V;A272V;I310L;F313M) does not contain a mutation at position 436. Third we hypothesize that tight packing in the binding pocket may lead to lower EC_{50} s. The docking results for I268V;A272V;I310L;F313M with LG335 show that the methionine and leucine side chains pack tightly against the propyl group of LG335, which may result in tighter binding and consequently a lower EC_{50} s.

In the absence of functional data, chemical complementation may be used to test more hypotheses about the function of particular residues than would be possible through site directed mutagenesis. By making a library of changes at a single site, additional information could be obtained about the importance of side chain size, polarity, and charge over just the traditional mutation to alanine that is often used to explore single residue importance. In the absence of structural information, it is possible to make large libraries using error prone PCR or gene shuffling. Chemical complementation could also be used to select active variants from these types of libraries.

Example 5: Increasing the Sensitivity of Chemical Complementation with ACTR.

To increase the sensitivity of chemical complementation, an adapter protein was introduced to link the mammalian nuclear receptor function to the yeast transcription apparatus, thereby overcoming the evolutionary divergence between mammalian cells and yeast. The human nuclear receptor coactivator ACTR was fused to the yeast Gal4 activation domain. This plasmid, pGAD10BACTR, expresses the ACTR:GAD fusion protein

and contains a leucine marker. This plasmid was co-transformed into yeast with the plasmid pGBDRXR, which expresses the Gal4 DNA binding domain (DBD) fused to the RXR ligand binding domain (GBD:RXR) and contains a tryptophan marker. Transformants were selected on SC -Leu-Trp plates, and
5 were streaked onto adenine selective plates (SC -Ade) containing 10^{-5} M 9cRA, a known ligand for RXR (Figure 5G).. Yeast containing just the pGBDRXR plasmid, the pGAD10BAACTR plasmid, a plasmid with just the Gal4DBD (pGBDMT), and a plasmid containing the Gal4 holo protein (pGBT9Gal4) were also streaked onto these plates as controls.

10 After two days of incubation, growth occurs on the sector of the plate containing ACTR:GAD with GBD:RXR and on the sector of the plate with Gal4; whereas no growth occurs on the sector of the plate with GBD:RXR alone (Figure 5G). The growth density produced by GBD:RXR and ACTR:GAD is the same as the growth produced by the holo Gal4.

15 Importantly, GBD:RXR and ACTR:GAD produced no growth on plates without 9cRA.

Previous findings showed no growth was observed with RXR at 9cRA concentrations lower than 10^{-5} M. To determine if the sensitivity of our system had increased with the introduction of the adapter fusion
20 protein, a dose response was performed on adenine selective plates (SC -Ade) containing ligand concentrations ranging from 10^{-5} M to 10^{-9} M. After two days of incubation, a clear dose response occurs on the plates (Figure 5). Without ligand, growth occurs only on the Gal4 sector of the plate, as expected. At concentrations as low as 10^{-8} M 9cRA, ligand-
25 activated growth occurs only on the sector of the plate containing both GBD:RXR with ACTR:GAD (Figure 5D). At concentrations of ligand above 10^{-8} M, higher density growth is observed on the sector of the plate containing GBD:RXR with ACTR:GAD. No growth occurs with GBD:RXR alone as expected. In summary, the introduction of the fusion protein
30 ACTR:GAD increases the sensitivity of chemical complementation. Growth occurs on adenine selective plates with 9cRA after two days of incubation (Figure 5). Ligand-activated growth is observed at 9cRA

concentrations as low as 10^{-8} M 9cRA. With chemical complementation, an approximate EC_{50} value between 10^{-8} M and 10^{-7} M for wild-type RXR and 9cRA, which is comparable to the EC_{50} value measured for wild-type RXR in mammalian cell assays ($\sim 10^{-7}$ M) (Figure 5). The growth density

and rate with the ACTR:GAD fusion protein is comparable to Gal4 activated growth. The same results were obtained on adenine selective plates (SC -Ade-Trp and SC -Ade-Leu-Trp) and on histidine selective plates (data not shown). In summary, introducing an adapter fusion protein of the human coactivator with the Gal4 activation domain increases the sensitivity of chemical complementation 1000-fold, making this system more efficient for analysis of protein/ligand interactions.

Example 6: Increasing Sensitivity of Chemical Complementation using SRC-1

Another RXR coactivator was tested to increase the sensitivity of chemical complementation. Residues 54 to 1442 of the human nuclear receptor coactivator, SRC-1, were fused to the Gal4 activation domain to construct the plasmid pGAD10BASRC1. This plasmid, which expresses SRC1 :GAD in yeast and contains a leucine marker was transformed with GBD:RXR; transformants selected from SC -Leu-Trp were streaked onto adenine selective plates (SC -Ade) with various concentrations of 9cRA (Figure 6). Ligand-activated growth is observed only in the sector of the plate containing both GBD:RXR with SRC1:GAD, and the same trend is observed with SRC-1 as the ACTR coactivator (Figure 6).

To verify that the increased sensitivity is from specific interactions between the coactivator and the active conformation of the receptor, a series of further controls was devised. pGAD10, a plasmid containing the Gal4 activation domain (GAD) without a coactivator domain was cotransformed with pGBDRXR. The plasmid was also transformed alone. pGAD10BAACTR, pGAD10BASRC1, pGBT9Gal4, and pGBDMT were all transformed individually. These controls were streaked onto adenine selective plates (SC -Ade) with and without 9cRA. In the absence of ligand, only the entire Gal4 gene (pGBT9Gal4) grows as expected (data not shown). In the presence of

10⁻⁵ M 9cRA, growth occurs with the GBD:RXR with ACTR:GAD and GBD:RXR with SRC1:GAD. The Gal4 AD only (without the coactivator domain) with GBD:RXR displays no growth. These results verify that the increase in chemical complementation is specifically due to the interaction of the coactivator fusion protein with the ligand-bound nuclear receptor (data not shown).

Example 7: Chemical complementation and negative selection

Negative selection is the opposite of classical genetic complementation. Instead of allowing the microbe to survive, a functional gene kills the microbe; only cells containing non-functional genes survive and form colonies on selective plates. Negative selection is useful for finding mutations that disrupt the function of a protein.

For negative selection in yeast, others have generated yeast strains that contain Gal4 response elements (REs) fused to the *URA3* gene. The *URA3* gene codes for orotidine-5'-phosphate decarboxylase, an enzyme in the uracil biosynthetic pathway. This gene can be used for both positive and negative selection. For positive selection, yeast expressing this gene will survive in the absence of uracil in the media. For negative selection, uracil and 5-fluoroorotic acid (FOA) is added to the media. Expression of orotidine-5'-phosphate decarboxylase converts FOA to the toxin 5-fluorouracil, which kills the yeast. As used herein, the term "negative chemical complementation" refers to negative selection that occurs due to the presence of a small molecule.

Plasmids pGBDRXR and pGAD10BAACTR were individually transformed and co-transformed into MaV103. Transformants were streaked onto uracil selective plates (SC -Ura-Trp) with 9cRA for positive selection (data not shown). The same trend was seen with the ACTR:GAD with GBD:RXR in the MaV103 strain as seen previously with the PJ69-4A strain. The same transformants were streaked onto selective plates (SC -Leu-Trp) with FOA for negative chemical complementation. Varying concentrations of 9cRA were also added to the plates, ranging from 10⁻⁵ M to 10⁻⁸ M. In the absence of ligand (Figure 7B), yeast grow on the sector of the plate

containing ACTR:GAD with GBD:RXR as expected. This is expected because uracil is provided, and in the absence of ligand RXR maintains its inactive conformation, preventing ACTR:GAD from binding and transcription does not occur. Without expression of the *URA3* gene, 5-fluorouracil is not produced and the yeast survive. However, as the concentration of ligand increases (Figure 7B-7F), less growth occurs and at the highest concentration of ligand, 10^{-5} M, very little growth occurs. The small amount of growth that is observed is due to background growth associated with negative selection in this strain.

Negative chemical complementation is advantageous for engineering receptors for new small molecules for several reasons. First, mutant receptor libraries may contain constitutively active receptors or receptors that activate transcription in response to endogenous small molecules. These undesirable receptors can be removed from the library with negative selection. Second, in some cases it will be desirable to remove members of the library that activate in response to certain small molecules, e.g. the natural ligands. Negative chemical complementation will remove these members of the library. The remaining library can then be put through chemical complementation with the small molecule of interest. Third, for enzyme engineering negative chemical complementation can remove library members that produce a particular small molecule, e.g. an enantiomer of the compound of interest. The remaining mutant enzyme library can then be put through chemical complementation to find those capable of producing the small molecule of interest. Fourth, for drug discovery, chemical libraries can be efficiently evaluated for antagonists of nuclear receptors by their ability to allow the yeast to survive negative chemical complementation.

Example 8: Chemical complementation with RXR mutants.

Several RXR mutants previously tested in both mammalian cell assays and with chemical complementation in yeast (without the coactivator fusion protein) showed a general, but less than complete correlation. Without the coactivator fusion protein, ligand-activated growth was observed only with

wild-type RXR and the F439L mutant after five days of incubation; none of the other mutants showed ligand-activated growth. The variation in the transcription machinery could lead to the different patterns in activation. To test whether the adapter fusion protein could overcome the differences and show a more direct correlation, all the mutants in Table 3 were cloned into pGBD vectors and cotransformed into yeast with pGAD10BAACTR. Again, transformants were selected from SC -Leu-Trp plates and then streaked onto adenine selective plates (SC -Ade-Trp). These mutants were tested with 9cRA and LG335 (a near-drug, a synthetic compound structurally similar to an RXR agonist but that does not activate wild-type RXR) (Table 3).

The transcriptional activation patterns of these mutants in chemical complementation with the addition of ACTR:GAD was observed on dose response plates containing both 9cRA and the synthetic ligand, LG335 (Figure 8). On the plate without ligand, growth occurs on the sector of the plate containing Gal4, but growth also occurs on the sector of the plate with the two mutants F313I and F313I;F439L. This could be a result of the mutations causing a structural modification to the binding pocket that is favorable for the binding of an endogenous small molecule in yeast. At 10^{-5} M 9cRA, growth occurs on the sectors of the plate with the single mutants, C432G, Q275C, I268F, I310M, V342F, and F439L, as well as some of the triple mutants I310M;F313I;F439L and Q275C;F313I;V342F. As the concentration of ligand decreases, some mutants no longer show ligand-activated growth. At 10^{-7} M 9cRA, growth is observed with the F439L mutant as well as wild-type RXR (Figure 8). At the lowest concentration of ligand, 10^{-8} M 9cRA, growth is observed in the Gal4 and F313I sectors of the plates. For the synthetic ligand LG335, growth is observed with several of the single, double and triple mutants at 10^{-5} M (Figure 8). At lower concentrations of ligand, the single mutants do not show much growth. However, several of the double and triple mutants I310M;F313I;F439L, Q275C;F313I, and I310M;F313I display ligand-activated growth at 10^{-7} M LG335. At 10^{-8} M LG335, some growth is still observed in the I310M;F313I;F439L sector of the plate.

A correlation is apparent between yeast growth and transcriptional activation in mammalian cells when quantitating these results and comparing them with results from cell culture assays (Table 3). The I268F, Q275C, C432G, I310M, and I310M; F313I; F439L mutations which had previously not shown any growth with chemical complementation, grow with the ACTR:GAD fusion protein (Figure 8). The more direct correlation between chemical complementation and mammalian cell assays shows that the coactivator fusion protein (ACTR:GAD) serves to bridge millions of years of evolution by adapting mammalian nuclear receptor function to the yeast transcription machinery.

Definitions

As used herein, the term "polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. Thus, for instance, polynucleotides as used herein refers to, among others, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. The terms "nucleic acid," "nucleic acid sequence," or "oligonucleotide" also encompasses a polynucleotide as defined above.

In addition, polynucleotide as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide.

It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term polynucleotide as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including simple and complex cells, inter alia.

The term "oligonucleotide" refers to relatively short polynucleotides.

Typically the term refers to single-stranded deoxyribonucleotides, but it can refer as well to single-or double-stranded ribonucleotides, RNA:DNA hybrids and double-stranded DNAs, among other compounds containing multiple nucleotides
5 linked through phosphodiester bonds. The phosphodiester bonds are typically 5'-3' linkages between the deoxyribose or ribose sugars of adjacent nucleotides, which is the predominant mode of nucleotide coupling in natural DNA or RNA, respectively. The nucleotides of an oligonucleotide can be the naturally occurring ribonucleotides, rA, rC, rG and rU; deoxyribonucleotides, dA, dC, dG and dT; or
10 other compounds in which the backbone and/or the base moieties differ from the standard nucleotides of DNA and RNA.

The term "non-natural" means not typically found in nature including those items modified by man. Non-natural includes chemically modified subunits such as nucleotides as well as biopolymers having non-natural linkages, backbones,
15 or substitutions.

The term "non-natural backbone" means a covalent chemical linkage that couples together two or more nucleotides in a manner that is not identical to the naturally-occurring RNA or DNA phosphodiester backbones. Chemical deviations from the natural backbone can include, but are not limited to,
20 chemical modification of a single site on the natural backbone or the replacement of a component of the backbone with a completely different chemical group. Methylation of the O2' site on the ribose sugar is an example of a chemical difference from the natural backbone that would constitute a non-natural backbone. Replacement of the ribose sugar with a hexose sugar and/or
25 replacement of the phosphate group in DNA or RNA with a phosphorothioate group are also examples of non-natural backbones. Exemplary modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-
30 alkylene phosphonates, 5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, selenophosphates and borano-phosphates having normal 3'-5' linkages, 2'-5' linked analogs of these,

and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Representative oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof).

Some oligonucleotide backbones do not include a phosphorus atom therein and have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

Some embodiments synthesize or use oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular -CH₂-NH-O-CH₂-, -CH₂-N(CH₃)-O-CH₂- [known as a methylene (methylimino) or MMI backbone], -CH₂-O-N(CH₃)-CH₂-, -CH₂-N(CH₃)-N(CH₃)-CH₂- and -O-N(CH₃)-CH₂-CH₂- [wherein the native phosphodiester backbone is represented as -O-P-O-CH₂-] of the above referenced U.S. Pat. No. 5,489,677, and the amide backbones of the above referenced U.S. Pat. No. 5,602,240.

In other embodiments, the disclosed methods and compositions may comprise modified oligonucleotides containing one or more substituted sugar moieties. Other modified oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Particularly preferred are O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other oligonucleotides comprise one of the following at the 2' position: C₁ to

C₁₀ lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an
5 intercalator, a group for improving pharmacokinetic properties and other substituents having similar properties. Another modification includes 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al. (1995) *Helv. Chim. Acta*, , 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoethoxy, i.e.,
10 a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethyl-aminoethoxy-ethyl or 2'-DMAEOE), i.e., 2'-O-CH₂-O-CH₂-N(CH₃)₂.

Other modifications include 2'-methoxy (2'-O-CH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂), 2'-allyl (2'-CH₂-CH=CH₂), 2'-O-allyl (2'-O-CH₂-CH=CH₂)
15 and 2'-fluoro (2'-F). The 2'-modification may be in the arabino (up) position or ribo (down) position. An exemplary 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide.

20 Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar.

A further modification includes Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 3' or 4' carbon atom of the sugar ring thereby forming a bicyclic sugar moiety. The linkage is preferably a methylene (-CH₂-)_n
25 group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 or 2. LNAs and preparation thereof are described in U.S. Patent No. 6,268,490 and WO 99/14226.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or
30 "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-

propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine(1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g., 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in The Concise Encyclopedia of Polymer Science and Engineering, pages 858-859, Kroschwitz, J. I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., Angewandte Chemie, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y. S., Chapter 15, Antisense Research and Applications, pages 289-302, Crooke, S. T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases may be particularly useful for increasing the binding affinity of the oligomeric compounds of the disclosure. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2.degree. C. (Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., eds., Antisense Research and Applications, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

The terms "including", "such as", "for example" and the like are intended to refer to exemplary embodiments and not to limit the scope of the present disclosure.

The term "polypeptides" includes proteins and fragments thereof.

5 Polypeptides are disclosed herein as amino acid residue sequences. Those sequences are written left to right in the direction from the amino to the carboxy terminus. In accordance with standard nomenclature, amino acid residue sequences are denominated by either a three letter or a single letter code as indicated as follows: Alanine (Ala, A), Arginine (Arg, R), Asparagine (Asn, N),
10 Aspartic Acid (Asp, D), Cysteine (Cys, C), Glutamine (Gln, Q), Glutamic Acid (Glu, E), Glycine (Gly, G), Histidine (His, H), Isoleucine (Ile, I), Leucine (Leu, L), Lysine (Lys, K), Methionine (Met, M), Phenylalanine (Phe, F), Proline (Pro, P), Serine (Ser, S), Threonine (Thr, T), Tryptophan (Trp, W), Tyrosine (Tyr, Y), and Valine (Val, V).

15 "Variant" refers to a polypeptide or polynucleotide that differs from a reference polypeptide or polynucleotide, but retains essential properties. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in
20 many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more modifications (e.g., substitutions, additions, and/or deletions). A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polypeptide may be naturally occurring such as an allelic variant, or it may be a variant that is not
25 known to occur naturally.

Modifications and changes can be made in the structure of the polypeptides of in disclosure and still obtain a molecule having similar characteristics as the polypeptide (e.g., a conservative amino acid substitution). For example, certain amino acids can be substituted for other amino acids in a
30 sequence without appreciable loss of activity. Because it is the interactive capacity and nature of a polypeptide that defines that polypeptide's biological functional activity, certain amino acid sequence substitutions can be made in a polypeptide sequence and nevertheless obtain a polypeptide with like properties.

In making such changes, the hydropathic index of amino acids can be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a polypeptide is generally understood in the art. It is known that certain amino acids can be substituted for other amino acids

5 having a similar hydropathic index or score and still result in a polypeptide with similar biological activity. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics. Those indices are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cysteine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4);
10 threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

It is believed that the relative hydropathic character of the amino acid determines the secondary structure of the resultant polypeptide, which in turn
15 defines the interaction of the polypeptide with other molecules, such as enzymes, substrates, receptors, antibodies, antigens, and the like. It is known in the art that an amino acid can be substituted by another amino acid having a similar hydropathic index and still obtain a functionally equivalent polypeptide. In such changes, the substitution of amino acids whose hydropathic indices are
20 within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

Substitution of like amino acids can also be made on the basis of hydrophilicity, particularly, where the biological functional equivalent polypeptide or peptide thereby created is intended for use in immunological embodiments.
25 The following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 \pm 1); glutamate (+3.0 \pm 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); proline (-0.5 \pm 1); threonine (-0.4); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5);
30 tryptophan (-3.4). It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent polypeptide. In such changes, the substitution of amino acids whose hydrophilicity values are within \pm

2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

As outlined above, amino acid substitutions are generally based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those of skill in the art and include (original residue: exemplary substitution): (Ala: Gly, Ser), (Arg: Lys), (Asn: Gln, His), (Asp: Glu, Cys, Ser), (Gln: Asn), (Glu: Asp), (Gly: Ala), (His: Asn, Gln), (Ile: Leu, Val), (Leu: Ile, Val), (Lys: Arg), (Met: Leu, Tyr), (Ser: Thr), (Thr: Ser), (Tyr: Trp, Phe), and (Val: Ile, Leu). Embodiments of this disclosure thus contemplate functional or biological equivalents of a polypeptide as set forth above. In particular, embodiments of the polypeptides can include variants having about 50%, 60%, 70%, 80%, 90%, and 95% sequence identity to the polypeptide of interest.

"Identity," as known in the art, is a relationship between two or more polypeptide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods, including, but not limited to, those described in (Computational Molecular Biology, Lesk, A. M., Ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., Ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part I, Griffin, A. M., and Griffin, H. G., Eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., Eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., SIAM J Applied Math., 48: 1073 (1988).

Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. The percent identity between two sequences can be determined by using analysis software (i.e., Sequence Analysis Software Package of the Genetics Computer Group, Madison Wis.) that incorporates the Needleman and Wunsch, (J. Mol. Biol., 48:

443-453, 1970) algorithm (e.g., NBLAST, and XBLAST). The default parameters are used to determine the identity for the polypeptides of the present invention.

By way of example, a polypeptide sequence may be identical to the reference sequence, that is be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence
5 such that the % identity is less than 100%. Such alterations are selected from: at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence
10 or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in the reference polypeptide by the numerical percent of the respective percent
15 identity (divided by 100) and then subtracting that product from said total number of amino acids in the reference polypeptide.

"Operably linked" refers to a juxtaposition wherein the components are configured so as to perform their usual function. For example, control sequences or promoters operably linked to a coding sequence are capable of
20 effecting the expression of the coding sequence.

As used herein, the term "transfection" refers to the introduction of a nucleic acid sequence into the interior of a membrane enclosed space of a living cell, including introduction of the nucleic acid sequence into the cytosol of a cell as well as the interior space of a mitochondria, nucleus or chloroplast. The
25 nucleic acid may be in the form of naked DNA or RNA, associated with various proteins or the nucleic acid may be incorporated into a vector.

As used herein, the term "vector" is used in reference to a vehicle used to introduce a nucleic acid sequence into a cell. A viral vector is virus that has been modified to allow recombinant DNA sequences to be introduced into host
30 cells or cell organelles.



The term "selective agent" refers to a substance that is required for growth or for preventing growth of a cell or microorganism, for example cells or microorganisms that have been engineered to require a specific substance for growth or inhibit or reduce growth in the absence of a complementing factor.

Exemplary complementing factors include enzymes that degrade the selective agent, or enzymes that produce a selective agent. Generally, selective agents include, but are not limited to amino acids, antibiotics, nucleic acids, minerals, nutrients, etc. Selective media generally refers to culture media deficient in at least one substance, for example a selective agent, required for growth. The addition of a selective agent to selective media results in media sufficient for growth.

As used herein, the term "coregulator" refers to a transcription modulator.

It should be emphasized that the above-described embodiments of the present disclosure, particularly, any "preferred" embodiments, are merely possible examples of implementations, merely set forth for a clear understanding of the principles of the disclosed subject matter. Many variations and modifications may be made to the above-described embodiment(s) without departing substantially from the spirit and principles of the disclosure. All such modifications and variations are intended to be included herein within the scope of this disclosure and protected by the following claims.

Table 3: Activation of Transcription by Wild-Type RXR and Variant Receptors in CV-1 Cells and in Yeast Strain PJ69-4α with pGAD-ACTR Plasmid and pGBDRXR-Mutant Plasmids.

Ligand	 9cRA		 LG335	
	Activation in Mammalian Cells†	Activation in Yeast Cells‡	Activation in Mammalian Cells†	Activation in Yeast Cells‡
Wild-type	+++++	+++++	---	---
I268F	+++++	+++++	---	---
Q275C	+++++	+++++	---	---
L326F	++	+	---	---
C432G	+++++	+++++	---	---
L309V	+++	---	---	---
L436V	+	---	---	---
I310M	++++	++++	+	+
F313I	+++++	Constitutive	++++	Constitutive
V342F	++	++	---	---
F439L	+++++	+++++	+++	+++
Q275C:I310M	+++	---	+	---
Q275C:F313I	+++	---	+++++	+++++
I310M:F313I	++++	++	+++++	+++++
F313I:L426V	+	---	+++	+++++
F313I:F439L	+++++	Constitutive	+++++	Constitutive
Q275C:I310M:F313I	+	---	++++	+++++
Q275C:F313I:V342F	+++	+++	+++	+++++
Q275C:L436V:F439L	+++	+	+	---
I310M:F313I:F439L	+++++	+++	+++++	+++++

†Key: +++++ > 110% of wild-type activation by ligand; ++++ 90-110% of wild-type activation by ligand; +++ 70-90% of wild-type activation by ligand; ++ 50-70% of wild-type activation by ligand; + 30-50% of wild-type activation by ligand; + 10-30% of wild-type activation by ligand (estimated uncertainty one +).

‡Key: +++++ > 110% of pGBT9-Gal 4 activation by ligand; ++++ 90-110% of pGBT9-Gal 4 activation by ligand; +++ 70-90% of pGBT9-Gal 4 activation by ligand; ++ 50-70% of pGBT9-Gal 4 activation by ligand; + 30-50% of pGBT9-Gal 4 activation by ligand; + 10-30% of pGBT9-Gal 4 activation by ligand (estimated uncertainty one +).

--- = no growth

Table 3

Table 4: Nine of nine sequences randomly chosen from the unselected library all contain background sequences, not the designed sequences. Twelve of twelve sequences randomly chosen from the selected library (indicating that they are functional) all contain designed sequences. This result demonstrates that chemical complementation is very efficient at identifying functional receptors within a large collection of nonfunctional receptors.

Unselected Library						
	I268	A271	A272	I310	F313	L436
1		deleted		deleted		deleted
2		deleted		deleted		deleted
3	GTA (V)	CCT (P)	CCT (P)		TCG (S)	deleted
4		deleted		deleted		deleted
5		deleted		deleted		GCG (A)
6		deleted		deleted		deleted
7		deleted		deleted		deleted
8		deleted		deleted		deleted
9		deleted		deleted		TTC (F)

Selected Library						
	I268	A271	A272	I310	F313	L436
1	GTG (V)		GTG (V)	TTG (L)	ATG (M)	TTG
2	GTG (V)		GCA	GTG (V)	TCC (S)	TTG
3	CTA (L)	GCT	GCA	ATG (M)	GTG (V)	TTG
4	GCG (A)		GCA	TCC (S)	GTG (V)	TTC (F)
5	GCT (A)	GCT	GCA	GCC (A)	GCG (A)	TTC (F)
6	GCT (A)	GCT	GTT (V)	GCC (A)	GCG (A)	TTC (F)
7	CTT (L)	GCT	GCT	GTC (V)	ATC (I)	TTG
8	CTG (L)	GTG (V)	GCG	TTG (L)	TTG (L)	TTG
9	GTG (V)	GTG (V)	GCG	TTG (L)	GTG (V)	TTG
10	GTA (V)		GTG (V)	ATG (M)	TCC (S)	ATG (M)
11	GCG (A)	GCG	GCA	ATG (M)	GCG (A)	ACG (T)
12	GCG (A)	GCT	GCG	TCG (S)	GTC (A)	TTC (F)

Table 4

CLAIMS

We claim:

- 5 1. A method for identifying receptors, comprising:
 - (a) introducing a first polynucleotide encoding a receptor in to a cell, wherein the receptor comprises a ligand binding domain for a target ligand operably linked to a polynucleotide binding domain so that binding of the target ligand to the receptor activates transcription of a second polynucleotide
 - 10 complementing a selection agent;
 - (b) culturing the cell on the selective media in the presence of the target ligand, wherein growth of the cell indicates interaction of the receptor with the target ligand.
- 15 2. The method of claim 1, further comprising culturing the cell on selective media in the absence of the target ligand, wherein growth of the cell indicates the receptor constitutively activates transcription of the second polynucleotide.
- 20 3. A cell comprising:
 - (a) a recombinant nuclear receptor that induces expression of a
 - 20 first polynucleotide in response to interaction with a target substance, wherein expression of the first polynucleotide complements a selective agent; and
 - (b) an adapter fusion protein comprising a human coregulator domain operably linked to an activation domain, wherein the adapter fusion protein enhances transcription of the first polynucleotide induced by the
 - 25 recombinant nuclear receptor.
4. The cell of claim 3, wherein the cell is eukaryotic or prokaryotic.
5. The cell of claim 3, wherein the cell is a yeast cell.
6. A method for identifying enzymes comprising:
 - (a) introducing a first polynucleotide into a cell that is unable to
 - 30 grow on selective media, wherein the cell expresses a recombinant receptor polypeptide that activates transcription of a second polynucleotide in response to interaction of the recombinant receptor polypeptide with a target substance;
 - (b) culturing the cell on the selective media; and
 - (c) selecting the cell that grows on the selective media.

7. The method of claim 6, wherein the target substance is produced by a polypeptide encoded by the first polynucleotide.

8. The method of claim 6, wherein a single target substance induces a conformational change in the recombinant receptor to activate transcription.

5 9. The method of claim 6, wherein the target substance is unmodified.

10. The method of claim 6, wherein growth on the selective media indicates the first polynucleotide encodes a product that complements the selective media.

10 11. The method of claim 6, wherein the cell is a eukaryotic or prokaryotic cell.

12. The method of claim 6, wherein the selective media does not contain an amino acid necessary for survival.

13. The method of claim 12, wherein the amino acid is selected from the group consisting of histidine and alanine.

15 14. The method of claim 6, wherein the first polynucleotide encodes an enzyme that produces the target substance.

15. The method of claim 6, wherein the transformed cell further expresses an adaptor fusion protein.

16. The method of claim 7, wherein the adaptor fusion protein 16.

20 17. The method of claim 6, wherein the first polynucleotide encodes an engineered enzyme.

18. The method of claim 6, wherein the first polynucleotide encodes a naturally occurring enzyme.

25 19. The method of claim 6, comprises a human coactivator for transcription of the second polynucleotide.

20. A cell comprising:

(a) a recombinant nuclear receptor that induces transcription of a first polynucleotide in response to interaction with a target substance; and

30 (b) an adapter fusion protein comprising a human coactivator domain operably linked to an activation domain, wherein the adapter fusion protein enhances transcription of the first polynucleotide induced by the recombinant nuclear receptor.

21. The cell of claim 20, wherein the human coactivator domain is selected from the group consisting of SRC-1 and ACTR.

22. The cell of claim 20, wherein the cell is unable to grow on selective media.

23. A method for selecting cells comprising:

(a) introducing a first polynucleotide into a cell, wherein the cell expresses a recombinant receptor polypeptide that activates transcription of a second polynucleotide in response to interaction of the recombinant receptor polypeptide with a target substance;

(b) culturing the cell on selective media in the presence of a first selection agent; and

(c) selecting the cell that survives on the selective media in the presence of the selection agent, wherein expression of the second polynucleotide inhibits growth of the cell.

24. The method of claim 23, wherein the second polynucleotide encodes a cytotoxic polypeptide.

25. The method of claim 24, wherein the cytotoxic polypeptide comprises a proapoptotic polypeptide.

26. The method of claim 21, wherein the first selective agent comprises 5-fluoroorotic acid.

27. The method of claim 26, wherein the second polynucleotide encodes orotidine-5'-phosphate decarboxylase.

28. The method of claim 27, wherein the toxic substance comprises 5-fluorouracil.

29. A method for assembling an enzymatic pathway comprising:

(a) introducing a plurality of polynucleotides encoding enzymes having different substrates into a cell that is unable to grow on selective media, wherein the cell expresses a recombinant receptor polypeptide that activates transcription of a second polynucleotide in response to interaction of the recombinant receptor polypeptide with a target substance;

(b) culturing the cell on the selective media; and

(c) selecting the cell that grows on the selective media, wherein growth of a cell on the selective media indicates that the plurality of polynucleotides encode enzymes for producing products that complement the selective media.

30. The method of claim 31, wherein the product of one of the enzymes is the substrate of another of the enzymes.

31. A method for identifying receptors, comprising:

- 5 (a) introducing a first polynucleotide encoding a receptor in to a cell, wherein the receptor comprises a ligand binding domain for a target ligand operably linked to a response element so that binding of the target ligand to the receptor activates transcription of a second polynucleotide complementing a selection agent;
- 10 (b) culturing the cell on the selective media in the presence of the target ligand, wherein growth of the cell indicates interaction of the receptor with the target ligand.

THIS PAGE BLANK (USPTO)

1/13




FIG. 1

2/13




FIG. 2

3/13




FIG. 3

4/13






FIG 4

5/13



6/13




FIG. 7

7/13





FIG. 8

8/13




FIG. 9

9/13




FIG. 10




FIG. 11

10/13




FIG. 12

11/13




FIG. 13




FIG. 14




FIG. 15

12/13






FIG. 16

13/13.



THIS PAGE BLANK (CONT.)

SEQUENCE LISTING

APPROPRIATE COPY 17 MAY 2006

<110> Georgia Tech Research Corporation
Donald, Doyle F.
Bahareh, Azizi
Lauren, Schwimmer J.

<120> ENGINEERING ENZYMES THROUGH GENETIC SELECTION

<130> 820701-2810

<150> 60/520,754

<151> 2003-11-17

<150> 60/520,813

<151> 2003-11-17

<160> 18

<170> PatentIn version 3.3

<210> 1

<211> 17

<212> DNA

<213> artificial sequence

<220>

<223> Cassette 1, F

<400> 1

cggaatttcc catgggc

17

<210> 2

<211> 37

<212> DNA

<213> artificial sequence

<220>

<223> BPF

<400> 2

ctcgccgaac gaccgggtca ccgcatgcc ctagtg

37

<210> 3

<211> 36

<212> DNA

<213> artificial sequence

<220>

<223> BPr

<400> 3

ccgcttgcc cactccacta gtggcatgcg gtgacc

36

<210> 4

<211> 37
<212> DNA
<213> artificial sequence

<220>
<223> cassette 2, BPf, BPr, SEf

<400> 4
cgggcaggct ggaatgagct cctcgacgga attctcc
37

<210> 5
<211> 36
<212> DNA
<213> artificial sequence

<220>
<223> SEr

<400> 5
cagcccgggtg gccaggagaa ttccgtcgag gagctc
36

<210> 6
<211> 40
<212> DNA
<213> artificial sequence

<220>
<223> cassette 3, SEf, SEr, AMf

<400> 6
ctctgcgctc catcgggctt aagtgccac caattgacac
40

<210> 7
<211> 46
<212> DNA
<213> artificial sequence

<220>
<223> AMr

<400> 7
ctccagcatc tccataagga aggtgtcaat tgggtgggcac ttaagc
46

<210> 8
<211> 17
<212> DNA
<213> artificial sequence

<220>
<223> cassette 4, AMf, AMr, and R

<400> 8

caaaggatgg gccgcag
17

<210> 9
<211> 46
<212> DNA
<213> artificial sequence

<220>
<223> oligos BP1

<400> 9
ggcaaacatg gggctgaacc ccagctcgcc gaacgaccg gtcacc
46

<210> 10
<211> 66
<212> DNA
<213> artificial sequence

<220>
<223> BP2

<220>
<221> misc_feature
<222> (33)..(33)
<223> A, C or T

<220>
<221> misc_feature
<222> (34)..(34)
<223> A or G

<220>
<221> misc_feature
<222> (35)..(35)
<223> C or G

<220>
<221> misc_feature
<222> (36)..(36)
<223> A, C or T

<220>
<221> misc_feature
<222> (37)..(37)
<223> A or G

<220>
<221> misc_feature
<222> (38)..(38)
<223> C or G

<220>
<221> misc_feature
<222> (45)..(45)
<223> A, C or T

<220>
<221> misc_feature
<222> (46)..(46)
<223> A or G

<220>
<221> misc_feature
<222> (47)..(47)
<223> C or G

<400> 10
gccactcca ctagtgtgaa aagctgtttg tcnnnnnntt ggcanngttg gtgaccgggt
60

cgttcg
66

<210> 11
<211> 48
<212> DNA
<213> artificial sequence

<220>
<223> BP3

<400> 11
cttttcacac tagtggagtg ggccaagcgg atcccacact tctcagag
48

<210> 12
<211> 28
<212> DNA
<213> artificial sequence

<220>
<223> BP4

<400> 12
ggggcagctc tgagaagtgt gggatccg
28

<210> 13
<211> 48
<212> DNA
<213> artificial sequence

<220>
<223> SE1

<220>
<221> misc_feature
<222> (22)..(22)
<223> A, G or T

<220>
<221> misc_feature
<222> (23)..(23)

<223> C or T

<220>

<221> misc_feature

<222> (24)..(24)

<223> G or C

<220>

<221> misc_feature

<222> (31)..(31)

<223> A, G or T

<220>

<221> misc_feature

<222> (32)..(32)

<223> C or T

<220>

<221> misc_feature

<222> (33)..(33)

<223> G or C

<400> 13

gcaggctgga atgagctcct cnnngcctcc nnntcccacc gctccatc
48

<210> 14

<211> 46

<212> DNA

<213> artificial sequence

<220>

<223> SE2

<400> 14

ccggtggcca ggagaattcc gtccttcacg gcgatggagc ggtggg
46

<210> 15

<211> 63

<212> DNA

<213> artificial sequence

<220>

<223> AM1

<220>

<221> misc_feature

<222> (38)..(38)

<223> A, G or T

<220>

<221> misc_feature

<222> (39)..(39)

<223> C or T

<220>

<221> misc_feature

<222> (40) .. (40)
<223> G or C

<400> 15
ggctctgcgc tccatcgggc ttaagtgcct ggaacatnnn ttccttcttc aagctcatcg
60

ggg
63

<210> 16
<211> 51
<212> DNA
<213> artificial sequence

<220>
<223> AM2

<400> 16
gcatctcaat aaggaaggtg tcaattgtgt gtccccgatg agcttgaaga a
51

<210> 17
<211> 12414
<212> DNA
<213> artificial sequence

<220>
<223> pGAD10BA seq 020803

<400> 17
gcttgcatgc aacttctttt cttttttttt cttttctctc tccccggtg ttgtctcacc
60

atatccgcaa tgacaaaaaa aatgatggaa gacactaaag gaaaaaatta acgacaaaga
120

cagcaccaac agatgtcggt gttccagagc tgatgagggg tatcttcgaa cacacgaaac
180

tttttcttc cttcattcac gcacactact ctctaattgag caacggtata cggccttctt
240

tccagttact tgaatttgaa ataaaaaaag tttgccgctt tgctatcaag tataaataga
300

cctgcaatta ttaatctttt gtttctctgt cattgttctc gtccctttc ttccttggtt
360

ctttttctgc acaatatttc aagctatacc aagcatacaa tcaactccaa gctttgcaaa
420

gatggataaa gcggaattaa ttccccgagcc tccaaaaaag aagagaaagg tcgaattggg
480

taccgcccgc aattttaatc aaagtgggaa tattgctgat agctcattgt ccttcacttt
540

cactaacagt agcaacggtc cgaacctcat aacaactcaa acaaattctc aagcgctttc
600

acaaccaatt gcctcctcta acgttcatga taacttcatg aataatgaaa tcacggctag
660

taaaattgat gatggtaata attcaaaacc actgtcacct ggttggacgg accaaactgc
720

gtataacgcg tttggaatca ctacagggat gtttaatacc actacaatgg atgatgtata
780

taactatcta ttcgatgatg aagatacccc accaaaacca aaaaaagaga tctttatgag
840

tggattagga gaaaacttgg atccactggc cagtgtattca cgaaaacgca aattgccatg
900

tgatactcca ggacaaggtc ttacctgcag tggtgaaaaa cggagacggg agcaggaaaag
960

taaatatatt gaagaattgg ctgagctgat atctgccaat cttagtata ttgacaattt
1020

caatgtcaaa ccagataaat gtgcgatttt aaaggaaaca gtaagacaga tacgtcaaat
1080

aaaagagcaa ggaaaaacta tttccaatga tgatgatgtt caaaaagccg atgtatcttc
1140

tacagggcag ggagttattg ataaagactc cttaggaccg cttttacttc aggcattgga
1200

tggtttccta tttgtggtga atcgagacgg aaacattgta tttgtatcag aaaatgtcac
1260

acaatacctg caatataagc aagaggacct ggttaacaca agtggtttaca atatcttaca
1320

tgaagaagac agaaaggatt ttcttaagaa tttacaaaaa tctacagtta atggagtttc
1380

ctggacaaat gagacccaaa gacaaaaaag ccatacattt aattgccgta tgttgatgaa
1440

aacaccacat gatattctgg aagacataaa cgccagtcct gaaatgcgcc agagatatga
1500

aacaatgcag tgctttgccc tgtctcagcc acgagctatg atggaggaag ggggaagattt
1560

gcaatcttgt atgatctgtg tggcacgccg cattactaca ggagaaagaa catttccatc
1620

aaacctgag agctttatta ccagacatga tctttcagga aaggttgtca atatagatac
1680

aaattcactg agatcctcca tgaggcctgg ctttgaagat ataatccgaa ggtgtattca
1740

gagatttttt agtctaaatg atgggcagtc atgggccag aaacgtcact atcaagaagt
1800

taccagtgat gggatatttt ccccaacagc ttatcttaat ggccatgcag aaacccagtc
1860

atatcgattc tcgttggtcg atggaactat agtgactgca cagacaaaaa gcaaactctt
1920

ccgaaatcct gtaacaaatg atcgacatgg ctttgtctca acccacttcc ttcagagaga
1980

acagaatgga tatagaccaa acccaaatcc tgttgacaa gggattagac cacctatggc
2040

tggatgcaac agttcggtag gcggcatgag tatgtcgcca aaccaaggct tacagatgcc
2100

gagcagcagg gcctatggct tggcagaccc tagcaccaca gggcagatga gtggagctag
2160

gtatgggggt tccagtaaca tagcttcatt gaccctggg ccaggcatgc aatcaccatc
2220

ttcctaccag aacaacaact ataggctcaa catgagtagc cccccacatg ggagtcctgg
2280

tcttgcccca aaccagcaga atatcatgat ttctcctcgt aatcgtggga gtccaaagat
2340

agcctcacat cagttttctc ctgttgacgg tgtgcactct cccatggcat cttctggcaa
2400

tactgggaac cacagctttt ccagcagctc tctcagtgcc ctgcaagcca tcagtgaagg
2460

tgtggggact tcccttttat ctactctgtc atcaccaggc cccaaattgg ataactctcc
2520

caatatgaat attacccaac caagtaaagt aagcaatcag gattccaaga gtccctctggg
2580

cttttattgc gacaaaaatc cagtggagag ttcaatgtgt cagtcaaata gcagagatca
2640

cctcagtgc aaagaaagta aggagagcag tgttgagggg gcagagaatc aaaggggtcc
2700

tttgaaagc aaaggtcata aaaaattact gcagttactt acctgttctt ctgatgaccg
2760

gggtcattcc tccttgacca actccccct agattcaagt tgtaaagaat cttctgttag
2820

tgtcaccagc cctctggag tctcctctc tacatctgga ggagtatcct ctacatccaa
2880

tatgcattgg tcaactgttac aagagaagca ccgattttg cacaagttgc tgcagaatgg
2940

gaattcacca gctgaggtag ccaagattac tgcagaagcc actgggaaag acaccagcag
3000

tataacttct tgtggggacg gaaatgttgt caagcaggag cagctaagtc ctaagaagaa
3060

ggagaataat gcacttctta gatacctgct ggacagggat gatcctagtg atgcactctc
3120

taaagaacta cagccccaag tggaaggagt ggataataaa atgagtcagt gcaccagctc
3180

caccattcct agctcaagtc aagagaaaga ccctaaaatt aagacagaga caagtgaaga
3240

gggatctgga gacttgata atctagatgc tattcttggt gatctgacta gttctgactt
3300

ttacaataat tccatatact caaatggtag tcatctgggg actaagcaac aggtgtttca
3360

aggaactaat tctctgggtt tgaaaagtgc acagtctgtg cagtctatcc gtcctccata
3420

taaccgagca gtgtctctgg atagccctgt ttctgttggc tcaagtcctc cagtaaaaaa
3480

tatcagtgc tccccatgt taccaaagca acccatgttg ggtgggaatc caagaatgat
3540

ggatagtcag gaaaattatg gctcaagtat gggagactgg ggcttaccaa actcaaaggc
3600

cggcagaatg gaacctatga attcaaactc catgggaaga ccaggaggag attataatac
3660

ttctttaccc agacctgcac tgggtggctc tattcccaca ttgctcttc ggtctaatag
3720

cataccaggt gcgagaccag tattgcaaca gcagcagcag atgcttcaaa tgaggcctgg
3780

tgaaatcccc atgggaatgg gggctaattc ctatggccaa gcagcagcat ctaaccaact
3840

gggttcctgg cccgatggca tgttgtccat ggaacaagtt tctcatggca ctcaaaatag
3900

gcctcttctt aggaattccc tggatgatct tgttgggcca ccttccaacc tggaaggcca
3960

gagtgcgaa agagcattat tggaccagct gcacactctt ctcagcaaca cagatgccac
4020

aggcctggaa gaaattgaca gagctttggg cattcctgaa cttgtcaatc agggacaggc
4080

attagagccc aaacaggatg ctttccaagg ccaagaagca gcagtaatga tggatcagaa
4140

ggcaggatta tatggacaga catacccagc acaggggcct ccaatgcaag gaggctttca
4200

tcttcagggga caatcaccat cttttaactc tatgatgaat cagatgaacc agcaaggcaa
4260

ttttcctctc caaggaatgc acccacgagc caacatcatg agaccccgga caaacacccc
4320

caagcaactt agaatgcagc ttcagcagag gctgcagggc cagcagtttt tgaatcagag
4380

ccgacaggca cttgaattga aaatggaaaa cctactgct ggtggtgctg cggatgatgag
4440

gcctatgatg cagccccagc agggttttct taatgctcaa atggtcgccc aacgcagcag
4500

agagctgcta agtcactact tccgacaaca gagggtggct atgatgatgc agcagcagca
4560

acagcagcag cagcagcagc agcagcagca acagcaacag caacagcaac agcagcaaca
4620

gcagcaaacc caggccttca gcccacctcc taatgtgact gcttccccca gcatggatgg
4680

gcttttggca ggacccacaa tgccacaagc tcctccgcaa cagtttccat atcaacaaa
4740

ttatggaatg ggacaacaac cagatccagc ctttggtcga gtgtctagtc ctcccaatgc
4800

aatgatgtcg tcaagaatgg gtccctccca gaatcccatg atgcaacacc cgcaggctgc
4860

atccatctat cagtcctcag aaatgaaggg ctggccatca ggaaatttgg ccaggaacag
4920

ctccttttcc cagcagcagt ttgcccacca ggggaatcct gcagtgtata gtatggtgca
4980

catgaatggc agcagtggc acatgggaca gatgaacatg aaccccatgc ccatgtctgg
5040

catgcctatg ggtcctgatc agaaatactg ctgacatctc tgcaccagga cctcttaagg
5100

aaaccactgt acaaatgaca ctgcactagg attattggga aggaatcatt gttccaggca
5160

tccatcttgg aagaaaggac cagctttgag ctccatcaag ggtattttta gtgatgtcat
5220

ttgagcagga ctggatttta agccgaaggg caatatctac gtgtttttcc cccctccttc
5280

tgctgtgtat catggtgttc aaaacagaaa tgttttttgg cattccacct cctagggata
5340

taattctgga gacatggagt gttactgac ataaaacttt tgtgtcactt ttttctgcct
5400

tgctagccaa aatctcttaa atacacgtag gtgggccaga gaacattgga agaatcaaga
5460

gagattagaa tatctggttt ctctagttgc agtattggac aaagagcata gtcccagcct
5520

tcagggtgtag tagttctgtg ttgacccttt gtccagtgga attggtgatt ctgaattgtc
5580

ctttactaat ggtgttgagt tgctctgtcc ctattatttg ccctaggctt tctcctaata
5640

aaggttttca tttgccattc atgtcctgta atacttcacc tccaggaact gtcattggatg
5700

tccaaatggc tttgcagaaa ggaaatgaga tgacagtatt taatcgcagc agtagcaaac
5760

ttttcacatg ctaatgtgca gctgagtgca ctttatttaa aaagaatgga taaatgcaat
5820

attcttgagg tcttgaggga atagtgaac acattcctgg tttttgcta cacttacgtg
5880

ttagacaaga actatgattt ttttttttaa agtactggtg tcaccctttg cctatatggt
5940

agagcaataa tgctttttta aaataaactt ctgaaaaccc aaggccaggt actgcattct
6000

gaatcagaat ctgcagtggt ttctgtgaat agattttttt gtaaataatga cctttaagat
6060

attgtattat gtaaaatatg tatatacctt tttttgtagg tcacaacaac tcatttttac
6120

agagtttgtg aagctaaata ttttaacattg ttgatttcag taagctgtgt ggtgaggcta
6180

ccagtggaag agacatccct tgacttttgt ggcctggggg aggggtagtg caccacagct
6240

tttccttccc cccccccag ccttagatgc ctgcgtcttt tcaatctctt aatctaaatg
6300

ctttttaag agattatttg ttttagatgta ggcattttta ttttttaaaa attcctctac
6360

cagaactaag cactttgtta atttgggggg aaagaataga tatggggaaa taaacttaaa
6420

aaaaaatcag gaatttaaaa aaaacgagca atttgaagag aatcttttgg attttaagca
6480

gtccgaaata atagcaattc atgggctgtg tgtgtgtgtg tatgtgtgtg tgtgtgtgtg
6540

tatgtttaat tatgttacct tttcatcccc tttaggagcg ttttcagatt ttggttcgta
6600

agacctgaat cccgcggccg ccccgggcgt agatactgaa aaaccccgca agttcacttc
6660

aactgtgcat cgtgcaccat ctcaatttct ttcatttata catcgttttg ccttctttta
6720

tgtaactata ctctctaaag tttcaatctt ggccatgtaa cctctgatct atagaatttt
6780

ttaaatgact agaattaatg cccatctttt ttttggacct aaattcttca tgaaaatata
6840

ttacgagggc ttattcagaa gctttggact tcttcgccag aggtttggtc aagtctccaa
6900

tcaaggttgt cggcttgtct accttgccag aaatttacga aaagatggaa aaggggtcaaa
6960

tcgttggtag atacgttggt gacacttcta aataagcgaa tttcttatga tttatgattt
7020

ttattattaa ataagttata aaaaaataa gtgtatacaa attttaaagt gactcttagg
7080

ttttaaaacg aaaattcttg ttcttgagta actctttcct gtaggtcagg ttgctttctc
7140

aggtatagca tgaggtcgct cttattgacc acacctctac cggcatgccc gaaatteccc
7200

taccctatga acatattcca ttttgtaatt tcgtgtcggt tctattatga atttcattta
7260

taaagtttat gtacaaatat cataaaaaaa gagaatcttt ttaagcaagg attttcttaa
7320

cttcttcggc gacagcatca ccgacttcgg tggacttggt ggaaccacct aaatcaccag
7380

ttctgatacc tgcattccaa acctttttaa ctgcatcttc aatggcctta ccttcttcag
7440

gcaagttcaa tgacaatttc aacatcattg cagcagacaa gatagtggcg ataggggtcaa
7500

ccttattctt tggcaaatct ggagcagaac cgtggcatgg ttcgtacaaa ccaaatgcgg
7560

tgttcttgct tggcaaagag gccaaaggac cagatggcaa caaacccaag gaacctggga
7620

taacggaggc ttcacggag atgatatcac caaacatggt gctggtgatt ataataccat
7680

ttaggtgggt tgggttctta actaggatca tggcggcaga atcaatcaat tgatgttgaa
7740

ccttcaatgt aggaaattcg ttcttgatgg ttctctccac agtttttctc cataatcttg
7800

aagaggccaa aacattagct ttatccaagg accaaatagg caatgggtggc tcatgttgta
7860

gggccatgaa agcggccatt cttgtgattc ttgacactc tggaacgggtg tattgttcac
7920

tatcccaagc gacaccatca ccacgtctt cctttctctt accaaagtaa atacctccca
7980

ctaattctct gacaacaacg aagtcagtag ctttagcaaa ttgtggcttg attggagata
8040

agtctaaaag agagtcggat gcaaagttac atggtcttaa gttggcgtac aattgaagtt
8100

ctttacggat ttttagtaaa ccttggtcag gtctaacact acctgtacct catttaggac
8160

caccacagc acctaacaaa acggcatcaa ccttcttgga ggcttcacgc gcctcatctg
8220

gaagtgggac acctgtagca tcgatagcag caccaccaat taaatgattt tcgaaatcga
8280

acttgacatt ggaacgaaca tcagaaatag ctttaagaac cttaatggct tcggctgtga
8340

tttcttgacc aacgtggtca cctggcaaaa cgacgatctt cttaggggca gacattagaa
8400

tggtatatcc ttgaaatata tatatatatt gctgaaatgt aaaaggtaag aaaaggtaga
8460

aagtaagacg attgctaacc acctattgga aaaaacaata ggtccttaaa taatattgtc
8520

aacttcaagt attgtgatgc aagcatttag tcatgaacgc ttctctattc tatatgaaaa
8580

gccggttccg gcctctcacc ttctcttttt ctcccaattt ttcagttgaa aaaggatat
8640

gcgtcaggcg acctctgaaa ttaacaaaaa atttcagtc atcgaatttg attctgtgcg
8700

atagcgcccc tgtgtgttct cgttatgttg aggaaaaaaa taatggttgc taagagattc
8760

gaactcttgc atcttacgat acctgagtat tcccacagtt ggggatctcg actctagcta
8820

gaggatcaat tcgtaatcat ggtcatagct gtttctcttg tgaaattgtt atccgctcac
8880

aattccacac aacatagcag ccggaagcat aaagtgtaaa gcctgggggtg cctaattgagt
8940

gaggtaactc acattaattg cggtgcgctc actgcccgtt ttccagtcgg gaaacctgtc
9000

gtgccagctg gattaatgaa tcggccaacg cgcggggaga ggcgggttgc gtattgggag
9060

ctcttccgct tcctcgctca ctgactcgct gcgctcggtc gttcggctgc ggcgagcggc
9120

atcagctcac tcaaaggcgg taatacgggt atccacagaa tcaggggata acgcaggaaa
9180

gaacatgtga gcaaaaggcc agcaaaaggc caggaaccgt aaaaaggccg cgttgctggc
9240

gtttttccat aggctccgcc cccctgacga gcatacaaaa aatcgacgct caagtcagag
9300

gtggcgaaac ccgacaggac tataaagata ccaggcggtt cccctggaa gctccctcgt
9360

gcgctctcct gttccgaccc tgccgcttac cggatacctg tccgccttcc tcccttcggg
9420

aagcgtggcg ctttctcata gctcacgctg taggtatctc agttcgggtg aggtcgcttcg
9480

ctccaagctg ggctgtgtgc acgaaccccc cgttcagccc gaccgctgcg ccttatccgg
9540

taactatcgt cttgagtcca acccggttaag acacgactta tcgccactgg cagcagccac
9600

tggtaacagg attagcagag cgaggtatgt aggcgggtgct acagagttct tgaagtgggtg
9660

gcctaactac ggctacacta gaaggacagt atttggtatc tgcgctctgc tgaagccagt
9720

taccttcgga aaaagagttg gtagctcttg atccggcaaaa caaaccaccg ctggtagcgg
9780

tgggtttttt gtttgcaagc agcagattac gcgcagaaaa aaaggatctc aagaagatcc
9840

tttgatcttt tctacggggt ctgacgctca gtggaacgaa aactcacgtt aagggatttt
9900

ggtcatgaga ttatcaaaaa ggatcttcac ctgatacctt ttaaattaaa aatgaagttt
9960

taaatcaatc taaagtatat atgagtaaac ttggtctgac agttaccaat gcttaatcag
10020

tgaggcacct atctcagcga tctgtctatt tcgttcaccc atagttgcct gactccccgt
10080

cggtgtagata actacgatac gggaggggctt accatctggc cccagtgcgt caatgatacc
10140

gcgagacca cgctcaccgg ctccagatTT atcagcaata aaccagccag ccggaagggc
10200

cgagcgcaga agtggctctg caactttatc cgcctccatc cagtctatta attgttgccg
10260

ggaagctaga gtaagtagtt cgccagttaa tagtttgccg aacgttggtg ccattgctac
10320

aggcatcgtg gtgtcacgct cgtcgtttgg tatggcttca ttcagctccg gttcccaacg
10380

atcaaggcga gttacatgat ccccatggt gtgcaaaaaa gcggttagct ccttcggtcc
10440

tccgatcgtt gtcagaagta agttggccgc agtggtatca ctcatggtta tggcagcact
10500

gcataattct cttactgtca tgccatccgt aagatgcttt tctgtgactg gtgagtactc
10560

aaccaagtca ttctgagaat agtgtatgcg gcgaccgagt tgctcttgcc cggcgctcaat
10620

acgggataat accgcgccac atagcagaac tttaaaagtg ctcatcattg gaaaacgttc
10680

ttcggggcga aaactctcaa ggatcttacc gctggtgaga tccagttcga tgtaaccac
10740

tcgtgcaccc aactgatctt cagcatcttt tactttcacc agcgtttctg ggtgagcaaa
10800

aacaggaagg caaaatgccg caaaaaaggg aataaggcg acacggaaat gttgaatact
10860

catactcttc ctttttcaat attattgaag catttatcag ggttattgtc tcatgagcgg
10920

atacatattt gaatgtattt agaaaaataa acaaataggg gtccgcgca catttccccg
10980

aaaagtgcc cctgacgtct aagaaaccat tattatcatg acattaacct ataaaaatag
11040

gcgtatcacg aggcccttcc gtctcgcgcg ttccgggtgat gacggtgaaa acctctgaca
11100

catgcagctc ccggagacgg tcacagcttg tctgtaagcg gatgccggga gcagacaagc
11160

ccgtcagggc gcgtcagcgg gtgttgccgg gtgtcggggc tggcttaact atgcggcatc
11220

agagcagatt gtactgagag tgcaccataa cgcatttaag cataaacacg cactatgccg
11280

ttcttctcat gtatatatat atacaggcaa cgcgcagata taggtgcgac gtgaacagtg
11340

agctgtatgt ggcagctcg cgttgcattt tcggaagcgc tcgttttcgg aaacgctttg
11400

aagttcctat tccgaagttc ctattctcta gctagaaagt ataggaactt cagagcgctt
11460

ttgaaaacca aaagcgctct gaagacgcac tttcaaaaaa ccaaaaacgc accggactgt
11520

aacgagctac taaaatattg cgaataccgc ttccacaaac attgctcaaa agtatctctt
11580

tgctatatat ctctgtgcta tatccctata taacctaccc atccaccttt cgctccttga
11640

acttgcatct aaactcgacc tctacatttt ttatgtttat ctctagtatt actctttaga
11700

caaaaaaatt gtagtaagaa ctattcatag agtgaatcga aaacaatacg aaaatgtaaa
11760

catttcctat acgtagtata tagagacaaa atagaagaaa ccgttcataa tttcttgacc
11820

aatgaagaat catcaacgct atcactttct gttcacaaag tatgcgcaat ccacatcggt
11880

atagaatata atcggggatg cctttatctt gaaaaaatgc acccgagct tcgctagtaa
11940

tcagtaaacg cgggaagtgg agtcaggctt tttttatgga agagaaaata gacaccaaag
12000

tagccttctt ctaaccttaa cggacctaca gtgcaaaaag ttatcaagag actgcattat
12060

agagcgca aaggagaaaa aaagtaatct aagatgcttt gttagaaaaa tagcgctctc
12120

gggatgcatt tttgtagaac aaaaaagaag tatagattct ttgttggtaa aatagcgctc
12180

tcgcgttgca tttctgttct gtaaaaatgc agctcagatt ctttgtttga aaaattagcg
12240

ctctcgctt gcatttttgt tttacaaaaa tgaagcacag attcttcggt ggtaaaatag
12300

cgctttcgcg ttgcatttct gttctgtaaa aatgcagctc agattctttg tttgaaaaat
12360

tagcgctctc gcgttgcat tttgttctac aaaatgaagc acagatgctt cgtt
12414

<210> 18

<211> 7254

<212> DNA

<213> artificial sequence

<220>

<223> pGBDRXRalpha

<400> 18

gcttgcatgc aacttctttt cttttttttt cttttctctc tccccggtg ttgtctcacc
60

atatccgcaa tgacaaaaaa aatgatggaa gacactaaag gaaaaaatta acgacaaaga
120

cagcaccaac agatgtcggt gtccagagc tgatgagggg tatcttcgaa cacacgaaac
180

tttttccttc cttcattcac gcacactact ctctaataag caacggtata cggccttcct
240

tccagttact tgaatttgaa ataaaaaaag tttgccgctt tgctatcaag tataaataga
300

cctgcaatta ttaatctttt gtttctctgt cattgttctc gttccctttc ttccttggtt
360

ctttttctgc acaatatttc aagctatacc aagcatataa tcaactccaa gcttgaagca
420

agcctcctga aagatgaagc tactgtcttc tatcgaacaa gcatgcgata tttgccgact
480

taaaaagctc aagtgtcca aagaaaaacc gaagtgcgcc aagtgtctga agaacaactg
540

ggagtgtcgc tactctccca aaacaaaaag gtctccgctg actagggcac atctgacaga
600

agtggaatca aggctagaaa gactggaaca gctatttcta ctgatttttc ctcgagaaga
660

ccttgacatg attttgaaaa tggattcttt acaggatata aaagcattgt taacaggatt
720

atttgtacaa gataatgtga ataaagatgc cgtcacagat agattggctt cagtggagac
780

tgatatgcct ctaacattga gacagcatag aataagtgcg acatcatcat cggaagagag
840

tagtaacaaa ggtcaaagac agttgactgt atcgccggaa tccccgggac agctgcattc
900

tcccatcagc accctgagct ccccatcaa cggcatgggc ccgcctttct cggtcacag
960

ctcccccatg ggccccact ccattgtcgt gccaccaca cccaccctgg gcttcagcac
1020

tggcagcccc cagctcagct cacctatgaa ccccgtcagc agcagcgagg acatcaagcc
1080

ccccctgggc ctcaatggcg tcctcaaggt ccccgccac ccctcaggaa acatggcttc
1140

cttcaccaag cacatctgcg ccatctgcgg ggaccgctcc tcaggcaagc actatggagt
1200

gtacagctgc gaggggtgca agggcttctt caagcggacg gtgcgcaagg acctgacct
1260

cacctgccgc gacaacaagg actgcctgat tgacaagcgg cagcgggaacc ggtgccagta
1320

ctgccgctac cagaagtgcc tggccatggg catgaagcgg gaagccgtgc aggaggagcg
1380

gcagcgtggc aaggaccgga acgagaaatga ggtggagtcg accagcagcg ccaacgagga
1440

catgccggtg gagaggatcc tggaggctga gctggccgtg gagcccaaga ccgagacct
1500

cgtggaggca aacatggggc tgaaccccag ctgccgaac gacctgtca ccaacattg
1560

ccaagcagcc gacaaacagc ttttcacctt ggtggagtgg gccaaagcga tcccacactt
1620

ctcagagctg cccctggacg accaggtcat cctgctgcgg gcaggctgga atgagctgct
1680

catcgctcc ttctcccacc gctccatcgc cgtgaaggac gggatcctcc tggccaccgg
1740

gctgcacgtc caccggaaca gcgcccacag cgcaggggtg ggcgccatct ttgacagggt
1800

gctgacggag cttgtgtcca agatgcggga catgcagatg gacaagacgg agctgggctg
1860

cctgcgcgcc atcgtcctct ttaacctga ctccaagggg ctctcgaacc cggccgaggt
1920

ggaggcgtg agggagaagg tctatgcgtc cttggaggcc tactgcaagc acaagtaccc
1980

agagcagccg ggaaggttcg ctaagctctt gctccgcctg ccggctctgc gctccatcg
2040

gctcaaatgc ctggaacatc tcttcttctt caagctcatc ggggacacac ccattgacac
2100

cttccttatg gagatgctgg aggcgccgca ccaaagact taggcctgcg ggcccatcct
2160

ttgtgcccac ccgttctggc caccctgcct ggacgccagc tgttcttctc agcctgagcc
2220

ctgtccctgc ccttctctgc ctggcctgtt tggactttgg ggcacagcct gtcactgctc
2280

tgcctaagag atgtgttgtc accctcctta tttctgttac tacttgtctg tggcccaggg
2340

cagtggctttt cctgagcagc agccttcgtg gcaagaacta gcgtgagccc agccaggcgc
2400

ctccccaccg ggctctcagg acgccctgcc acaccacgg ggcttgggcg actacagggg
2460

cttcggcccc agccctggag ctgcagccaa gctaattccg ggcgaatttc ttatgattta
2520

tgatttttat tattaaataa gttataaaaa aaataagtgt atacaaattt taaagtgact
2580

cttaggtttt aaaacgaaaa ttcttgttct tgagtaactc tttcctgtag gtcaggttgc
2640

tttctcaggt atagcatgag gtcgctctta ttgaccacac ctctaccggc atgccggcaa
2700

gtgcacaaac aataacttaaa taaatactac tcagtaataa cctatttctt agcatttttg
2760

acgaaatttg ctattttggt agagtctttt acaccatttg tctccacacc tccgcttaca
2820

tcaacaccaa taacgccatt taatctaagc gcatcaccaa cattttctgg cgtcagtcca
2880

ccagetaaca taaaatgtaa gctttcgggg ctctcttgcc ttccaacca gtcagaaatc
2940

gagttccaat ccaaaagttc acctgtccca cctgcttctg aatcaaacaa ggggaataaac
3000

gaatgaggtt tctgtgaagc tgcactgagt agtatgttgc agtcttttgg aaatacaggt
3060

cttttaataa ctggccttac tccaaagaca cttcgacgtg actcatcata caacgtcaga
3120

aaacctttat gctcagaaaa ttattgaccg aaaccgagga actcttggtt ttcttgccac
3180

gactcatctc catgcagttg gacgatatca atgccgtaat cattgaccag agccaaaaca
3240

tcctccttag gttgattacg aaacacgcca accaagtatt tcggagtgcc tgaactattt
3300

ttatatgctt ttacaagact tgaaattttc cttgcaataa ccgggtcaat tgttctcttt
3360

ctattgggca cacatataat acccagcaag tcagcatcgg aatctagagc acattctgcg
3420

gcctctgtgc tctgcaagcc gcaaactttc accaatggac cagaactacc tgtgaaatta
3480

ataacagaca tactccaagc tgcctttgtg tgcttaatac cgtatactca cgtgctcaat
3540

agtcaccaat gccctccctc ttggccctct ccttttcttt ttctgaccga attaatctgt
3600

aatcatgggc atagctgttt cctgtgtgaa attgttatcc gctcacaatt ccacacaaca
3660

tacgagccgg aagcataaag tgtaaagcct ggggtgccta atgagtgagg taactcacat
3720

taattgcgtt gcgctcactg cccgctttcc agtcgggaaa cctgtcgtgc cagctggatt
3780

aatgaatcgg ccaacgcgcg gggagaggcg gtttgcgat tgggcgctct tccgcttcc
3840

cgctcactga ctgcgtgcgc tcggtcgttc ggctgcggcg agcggatatca gctcactcaa
3900

aggcggtaat acggttatcc acagaatcag gggataacgc aggaaagaac atgtgagcaa
3960

aaggccagca aaaggccagg aaccgtaaaa aggcgcggtt gctggcggtt ttccataggc
4020

tccgcccccc tgacgagcat cacaaaaatc gacgctcaag tcagaggtgg cgaaaccgga
4080

caggactata aagataccag gcgtttcccc ctggaagctc cctcgtgcgc tctcctgttc
4140

cgaccctgcc gcttaccgga tacctgtccg cctttctccc ttccgggaagc gtggcgcttt
4200

ctcatagctc acgctgtagg tatctcagtt cgggtgtaggc cgttcgctcc aagctgggct
4260

gtgtgcacga acccccgtt cagcccgacc gctgcgcctt atccggtaac tatcgtcttg
4320

agtccaaccc ggtaagacac gacttatcgc cactggcagc agccactggc aacaggatta
4380

gcagagcgag gtatgtaggc ggtgctacag agttcttgaa gtggtggcct aactacggct
4440

acactagaag gacagtattt ggtatctgcg ctctgctgaa gccagttacc ttccgaaaaa
4500

gagttggtag ctcttgatcc ggcaaacaaa ccaccgctgg tagcgggtgg ttttttgtt
4560

gcaagcagca gattacgcgc agaaaaaaag gatctcaaga agatcctttg atcttttcta
4620

cggggtctga cgctcagtgg aacgaaaact cacgttaagg gattttggc atgagattat
4680

caaaaaggat cttcacctag atccttttaa attaaaaatg aagttttaaa tcaatctaaa
4740

gtatatatga gtaaacttgg tctgacagtt accaatgctt aatcagtgag gcacctatct
4800

cagcgatctg tctatttcgt tcatccatag ttgctgact ccccgtcgtg tagataacta
4860

cgatacggga gggcttacca tctggcccca gtgctgcaat gataccgcga gacccacgct
4920

caccggctcc agatttatca gcaataaacc agccagccgg aagggccgag cgcagaagtg
4980

gtcctgcaac tttatccgcc tccatccagt ctattaattg ttgccgggaa gctagagtaa
5040

gtagtccgcc agttaatagt ttgcgcaacg ttgttgccat tgctacaggc atcgtggtgt
5100

cacgctcgtc gtttggtatg gcttcattca gctccgggtc ccaacgatca aggcgagtta
5160

catgatcccc catgttgtgc aaaaaagcgg ttagctcctt cggtcctcgc atcgttgtca
5220

gaagtaagtt ggccgcagtg ttatcactca tggttatggc agcactgcat aattctctta
5280

ctgtcatgcc atccgtaaga tgcttttctg tgactggtga gtactcaacc aagtcattct
5340

gagaatagtg tatgcggcga ccgagttgct cttgcccggc gtcaataggg gataataccg
5400

cgccacatag cagaacttta aaagtgctca tcattggaaa acgttcttcg gggcgaaaac
5460

tctcaaggat cttaccgctg ttgagatcca gttcgatgta acccactcgt gcacccaact
5520

gatcttcagc atcttttact ttcaccagcg tttctgggtg agcaaaaaca ggaaggcaaa
5580

atgccgcaaa aaagggaata agggcgacac ggaaatgttg aatactcata ctcttccttt
5640

ttcaatatta ttgaagcatt tatcaggggtt attgtctcat gagcggatac atatttgaat
5700

gtatttagaa aaataaacia ataggggttc cgcgcacatt tccccgaaaa gtgccacctg
5760

acgtctaaga aaccattatt atcatgacat taacctataa aaataggcgt atcacgaggc
5820

cctttcgtct cgcgcgtttc ggtgatgacg gtgaaaacct ctgacacatg cagctcccgg
5880

agacggtcac agcttgtctg taagcggatg ccgggagcag acaagcccgt cagggcgcgt
5940

cagcgggtgt tggcgggtgt cggggctggc ttaactatgc ggcatcagag cagattgtac
6000

tgagagtgca ccataacgca tttaagcata aacacgcact atgccgttct tctcatgtat
6060

atatatatac aggcaacacg cagatatagg tgcgacgtga acagtgaagct gtatgtgcgc
6120

agctcgcggt gcattttcgg aagcgctcgt ttccggaaac gctttgaagt tcctattccg
6180

aagttcctat tctctagcta gaaagtatag gaacttcaga gcgcttttga aaacaaaaag
6240

cgctctgaag acgcactttc aaaaaaccaa aaacgcaccg gactgtaacg agctactaaa
6300

aaacttttgg ttttcgagag acttctgcgt gaaagttttt tggtttttgc gtggcctgac
6360

attgctcgat gatttatatt gcgaataccg cttccacaaa cattgctcaa aagtatctct
6420

ttgctatata tctctgtgct atatccctat ataacctacc catccacctt tcgctccttg
6480

aacttgcac taaactcgac ctctacattt tttatgttta tctctagtat tactctttag
6540

acaaaaaaat tgtagtaaga actattcata gagtgaatcg aaaacaatac gaaaatgtaa
6600

acatttccta tacgtagtat atagagacaa aatagaagaa accgttcata attttctgac
6660

caatgaagaa tcatcaacgc tatcactttc tgttcacaaa gtatgcgcaa tccacatcgg
6720

tatagaatat aatcggggat gcctttatct tgaaaaaatg caccgcgagc ttcgctagta
6780

atcagtaaac gcgggaagtg gagtcaggct ttttttatgg aagagaaaat agacacaaaa
6840

gtagccttct tctaacctta acggacctac agtgcaaaaa gttatcaaga gactgcatta
6900

tagagcgac aaaggagaaa aaaagtaatc taagatgctt tgtagaaaa atagcgctct
6960

cgggatgcat tttttagaaa caaaaaagaa gtatagattc tttgttggtta aaatagcgct
7020

ctcgcgttgc atttctgttc tgtaaaaatg cagctcagat tctttgtttg aaaaattagc
7080

gctctcgcgt tgcatttttg ttttacaaaa atgaagcaca gattcttcgt tggtaaaata
7140

gcgccttcgc gttgcatttc tgctctgtaa aaatgcagct cagattcttt gtttgaaaaa
7200

ttagcgctct cgcgttgcac ttttgctcta caaaatgaag cacagatgct tcgt
7254

THIS PAGE BLANK (USPTO)